

E

R

A

M

O

C

**Committee on Medical Aspects of Radiation
in the Environment (COMARE)**

FIFTEENTH REPORT

Radium contamination in the area around Dalgety Bay.

Chairman: Professor A Elliott

CONTENTS

Foreword

Chapter 1 Introduction

Chapter 2 Site history

Chapter 3 Radium, its use and disposal

Chapter 4 Contamination history

Chapter 5 Dosimetry and risk

Chapter 6 Cancer epidemiology

Chapter 7 Conceptual model and recent investigations

Chapter 8 Implications for other sites

Chapter 9 Conclusions

Chapter 10 Recommendations

References

Acknowledgements

Appendix A Glossary & Abbreviations

Appendix B COMARE Reports

Appendix C COMARE Membership & Writing Group Membership

Appendix D Declarations of Members' Interests

FOREWORD

- i. The Committee on Medical Aspects of Radiation in the Environment (COMARE) was established in November 1985 in response to the final recommendation of the report of the Independent Advisory Group chaired by Sir Douglas Black (Black, 1984). The terms of reference for COMARE are:

'to assess and advise Government and the devolved authorities on the health effects of natural and man-made radiation and to assess the adequacy of the available data and the need for further research'

- ii. In 28 years of providing advice to Government and the devolved authorities, COMARE has to date published 14 major reports (see Appendix B), in addition to numerous other statements and documents. These have been mainly related to exposure to naturally-occurring radionuclides such as radon and its progeny, or to man-made radiation, usually emitted by major nuclear installations. It has also published reports with a medical focus.
 - iii. COMARE has been considering radioactive contamination in the Dalgety Bay area since 1991 and has been involved in advising Scottish authorities since that date. Most advice has been channelled through the Scottish Environment Protection Agency (SEPA) since its inception in 1997.
-
- iv. In 2011, SEPA notified COMARE of a new series of finds of radioactive sources in higher numbers and containing greater amounts of radioactivity than in the preceding period and the Committee instituted appropriate investigations as detailed below. In November 2012, before the Committee had prepared a report, two internal working papers were released under a Freedom of Information request made to SEPA and COMARE. As a result of the subsequent press article, the Scottish Government requested that COMARE prepare an interim report by 31 December 2012 and this was submitted.
 - v. The aim of this COMARE report is to provide further information to the Scottish Government on the risks posed to the public by the contamination at Dalgety Bay, to examine the incidence of potentially related cancers and to recommend whether further actions to protect the public are required. Its methodology and approach may be applied to other sites where such contamination is found.

CHAPTER 1

INTRODUCTION

- 1.1 As noted above, COMARE has been considering radioactive contamination in the Dalgety Bay area since 1991 and has been involved in advising SEPA since its inception in 1997. A SEPA observer attends the meetings of the main COMARE committee, as does a Public Health England (PHE) observer, permitting detailed collaboration between the organisations. The Ministry of Defence (MoD) also has observer status, but has chosen not to attend committee meetings since July 2012.
- 1.2 Most of the early contamination found in the area of Dalgety Bay was in terrestrial locations. In these cases, each area affected was small and the contamination was immobile, easily delineated (i.e. a static problem) and amenable to remediation by excavation and removal.
- 1.3 The sources found from 2009 onwards were sited on the foreshore and appear to be evidence of a more dynamic (changing) situation. Under these circumstances, it is not possible to make accurate predictions about future hazard from past evidence.
- 1.4 In 2011, the most recent series of finds of radioactive sources located by SEPA was notified to COMARE. The Committee took the view that these finds represented a significant change to the previous situation, requiring more in depth investigation.
- 1.5 In November 2011, the Committee requested that the NHS Scotland Information Services Division (ISD) conduct an analysis of the incidence of cancers in the vicinity of Dalgety Bay, comparing local incidence rates with the national averages for the whole of Scotland. This was received by the committee in February 2012 and was copied also to NHS Fife.
- 1.6 The ISD report, covering the period 2000-2009, was considered by the Committee at its meeting in March 2012. Two types of cancer showed an increased incidence that was statistically significant – primary liver cancer and non-Hodgkin lymphoma (NHL). The Committee agreed that a further postcode analysis should be undertaken for these two cancers to determine whether there was any spatial pattern present that might indicate a link to the site of the contamination. It agreed also that Fife Health Board should be asked to conduct a case-note review of the liver cancer cases to double-check that these had been classified correctly.
- 1.7 The confidential report from Fife Health Board was received by the Chairman in July 2012 and the results of the postcode analysis for liver cancer were considered at the Committee's meeting in October 2012. The postcode analysis led to a request to ISD for further information and analyses for both liver cancer and NHL cases.
- 1.8 In November 2012, the ISD report and the postcode analysis paper were released under a Freedom of Information request made to SEPA and COMARE. As a result of the subsequent press article, the Scottish Government requested that COMARE prepare a report by 31 December 2012.
- 1.9 On several occasions, as far back as 1992, the Committee has requested information from the MoD on other current and former military sites where such contamination might be present. Through the Committee's efforts, the list has gradually lengthened and, most recently, the Committee's Contaminations Working Group sought information particularly on 'alienated' sites (those disposed of by the MoD) in September 2012, requesting the following information:
 - a) Are alienated sites monitored by the MoD?
 - b) Which alienated sites have been looked at in terms of monitoring or resurveying?

- c) What is the minimum detectable level of activity for the monitoring process?
- d) Does the MoD know of any potential pathways for human exposure at any of the sites?

To date, COMARE has not received an adequate response to these points.

CHAPTER 2

SITE HISTORY

- 2.1 Figure 2.1 shows the location of the site, which is on the north shore of the Firth of Forth, about 5 km to the east of the Forth road and rail bridges. It forms one of the Firth of Forth Sites of Special Scientific Interest (SSSI) and also is part of the Firth of Forth Ramsar sites (the Ramsar Convention is an international treaty for the conservation and sustainable utilisation of wetlands).



Figure 2.1 – Location of Dalgety Bay

Contains Ordnance Survey data © Crown copyright and database right 2013.

- 2.2 Figure 2.2 shows the locality, which now includes part of the town of Dalgety Bay, and delineates the area of interest. Information about ownership of the site in the following paragraphs has been taken from a SEPA report (SEPA, 2013) with permission.

- (HMS Merlin) was reduced to care and maintenance and was handed over to HMS Cochrane. The accommodation blocks were used as barracks for naval artificers being trained at Rosyth.
- 2.9 Parts of the area were used as agricultural and grazing land between the mid 1950s and 1960. The airfield was decommissioned during 1959 and was sold by the Ministry of Defence in 3 separate parts: 101 hectares to Moray Estates Development Company in 1962, 22 hectares to the Board of Trade in 1964 (see paragraph 2.13) and 27 hectares to Fife County Council in 1968 (see paragraph 2.14).
 - 2.10 In March 1960, Copthall Holdings (Dalgety) Development Limited was formed to build a new town at Dalgety Bay over the next 10 years, sponsored jointly with Moray Estates Development Company Limited. This was then passed to Copthall Holdings (Dalgety) Developments Limited in 1967 with the exception of the foreshore area which is still owned by Moray Estates Development Company Limited. Although Moray Estates Development Company still owns the foreshore, the sale to Copthall Holdings (Dalgety) Developments Limited does include the rights to the minerals and sand in the foreshore.
 - 2.11 Copthall Holdings (Dalgety) Developments Limited later changed its name to Donibristle Investments Limited and is now known as James Harrison Contracts Limited. The latter company still owns the land to the north and west of the land now owned by Dalgety Bay Sailing Club Limited, as well as the rights to the minerals and sand on the foreshore.
 - 2.12 The area was subsequently developed for residential and leisure uses. Outline planning permission for residential development was granted in May 1961. Groundworks for the infrastructure (roads, drainage etc) commenced in 1964 and the first houses were built in 1965 in the central area of the former airfield south of the former runway area. Development in the southern area of the former airfield started in the early 1970s and appears to have been completed by 1983.
 - 2.13 Land purchased by the Board of Trade was used to create Donibristle Industrial Estate. Plots or buildings were sold by the Board of Trade (subsequently the Scottish Development Agency and now Scottish Enterprise) to individual companies from 1968 onwards. The Scottish Development Agency had disposed of all land and property at Donibristle Industrial Estate by 1990.
 - 2.14 Land purchased by Fife County Council became Hillend Industrial Estate. Plots or buildings were sold by Fife County Council (subsequently Dunfermline District Council and Fife Regional Council) to individual companies starting in 1969 and continuing throughout the 1970s, 1980s and 1990s.
 - 2.15 Dalgety Bay Sailing Club was formed in 1971 and the current clubhouse built in 1975 with an extension added in 2000.
 - 2.16 Dalgety Bay Sailing Club was formally opened in 1972 on 2.6 acres of land leased from Donibristle Investments Limited at this time. The original clubhouse was a wooden garage located at the rear of where the existing top boat shed is now.
 - 2.17 The current clubhouse was completed in 1975. From the sailing club archives, it appears that the site was levelled and that material excavated from the foundations was used to create a mound against the masonry footings of the clubhouse on the southern elevation.
 - 2.18 A 1979 aerial photograph shows a second slipway in place to the west of the original slipway. The easternmost beach shows an increase in the number of boats being stored and, additionally, further storage is evident to the north in this area.

- 2.19 Evidence, confirmed by aerial photography, suggests that the rock armour (large quarried blocks) around the Headland and the Boat Park was placed and replenished throughout the 1980s (McPhail, 2011a). It appears that this material was added from the construction site of the Forth Road Bridge and is understood to be mainly smaller rocks and shale. The rock armour in front of the Headland area was augmented 8-10 years ago with larger rocks which also came from some additional construction work done in connection with the Forth Road Bridge. It is understood that the rock armour was emplaced by the Sailing Club and not by Fife Council (ibid).
- 2.20 The stonework of the New Harbour was restored in 1980 by members of the Sailing Club, no major earthworks being involved.
- 2.21 Dalgety Bay Sailing Club purchased two areas of land from James Harrison Contracts Limited, 1.26 acres of land in November 1984 with the pier or jetty known as New Harbour and sea walls thereon and also 2.85 acres of land in September 1985 bounded on the west by The Wynd.
- 2.22 Fife Council holds the responsibility for coastal defence. The current Fife Shoreline Management Plan (Fife Council, 2011) indicates that Fife Council have a general policy of 'Hold the Line' for currently defended sections of the coastline in the Dalgety Bay vicinity, but since the rock armour was not placed by the Council, it is not maintained by them. Information from Fife Council Transportation and Environmental Services Department indicates that they do not hold any records with respect to coastal defence works in the vicinity of Dalgety Bay.
- 2.23 Fife Council also has the responsibility for the maintenance of Ross Plantation, but has indicated that 'next to no works' within Ross Plantation have been carried out. More recently (since the 1990s) Fife Council, as a precautionary measure, has stopped any community planting or works by Council staff that might involve staff handling the soil, thus avoiding any contamination under their nails.

Coastal changes

- 2.24 The coastline of the area under consideration has changed over time and the changes have been assessed by means of aerial photographs.
- 2.25 Aerial photographs of the Dalgety Bay area were obtained from the Royal Commission on the Ancient and Historical Monuments of Scotland and analysed by the University of Stirling under a contract from SEPA (Sneddon, Tyler and Hunter, 2013). This dataset comprised 17 photographs taken between 1945 and 1990. The photographs were all taken on black and white panchromatic film, with exception of those collected in 1973.
- 2.26 The aerial photos were geocorrected to the British National Grid using Ordnance Survey (OS) map data (1:10,000 Raster product) by a third-order polynomial translation and nearest neighbour resampling. Initially, the 1990 aerial photo was geocorrected to the OS map using a large number of ground control points (GCPs; fixed points that could be accurately identified on both the OS map and the aerial photograph). The 1990 geocorrected aerial photograph was then used as the basis for the correction of the next photograph in the time sequence taken in 1986. This procedure was repeated with the remaining aerial photos geocorrected in reverse chronological order. This 'back stepping' approach minimised the variation in geolocational error with each time step.
- 2.27 The coastlines were identified by visual interpretation, photo contrast, brightness and gamma being modified through a range of interactive image stretches to assist with boundary identification. The coastlines were then superimposed over a 2009 Google MapsTM photo.

- 2.28 All photos in the period 1945 to 1950 showed a similar morphology for the bay. As the 1949 photo offered the best image quality, it was used as a baseline against which to compare changes that occur after 1950.
- 2.29 The coastline of the bay shown in the 1990 photo was fairly similar to that of the 2009 coastline, with overlap found for most of the coastline. Thus, it was assumed that the 1990 photo provides a relatively accurate representation of the current morphology of the coast line around Dalgety Bay. Figure 2.3 shows that there is some discrepancy between the 2009 site morphology and the 1990 morphology south of the study site; this discrepancy is caused by the expansion of the coastline in this area after 1990.




Figure 2.3 Shoreline contours for the west of the study area, the 'headland' area.

- 2.30 A substantial change in morphology occurs between 1949 and 1955 in the eastern part of the area (Figure 2.4), this change being characterised by a deformation to the south-eastern spit-like feature, suggesting that it accretes across the Bay in 1955. There is also a noticeable shift to the south of the Bay. The feature also appears to have above average shadowing, indicating an increased elevation relative to other features within the study area.
- 2.31 Between 1955 and 1959 the spit-like feature has been lost and the study area has been substantially infilled. This is established by the loss of exposed rock, which has been replaced by coarse sediment infill. This marks the time of most significant change with the extent of infilling being almost equivalent to the infilling reported for the entire study period.




Figure 2.4 Shoreline contours for the east of the study area, the Boatyard and 'Mud' Bay area.

- 2.32 Following the infilling period of 1955-1959, the coastline was found to vary slightly as accretion and erosion continued at the site for the following 30 years. Between 1959 and 1990, the coastline underwent minor changes with accretion in the north of the study site, particularly around the northern rock formation. In the south of the study site erosion occurs between 1959 and 1990 with the coastline retreating by several meters, before post 1990 expansion (Figure 2.3).
- 2.33 Having established the 1949 coastline as the optimal baseline and the 1990 as a reasonable representation of the current coastline, it was possible to estimate that an area of over 3,110 m² has been infilled over this period.
- 2.34 The headland (Figure 2.4) undergoes two primary changes, one post 1949 and the second post 1960. This headland interpretation uses higher quality photos as the headland does not benefit from a well-defined beach that is visible in all tidal conditions as is found at the bay study site.
- 2.35 The 1945-1949 headland coastlines demonstrate that there is little in the way of change between 1945 and 1949. Of importance however, is the "conical" shaped feature in the 1945 photo, an area marked as a refuse tip in later OS maps. This coastal section is seen to change over the subsequent years. Comparison of the 1949 and 1955 photos shows an expansion in the northern part of the headland occurring between 1949 and 1955. Over this time, the coastal part of the tip area is seen to change. There is a minor second expansion in headland post 1955 that occurs mid-way up the headland, as shown in the 1959 and 1960 headlands.
- 2.36 Sometime between 1960 and 1973, a sandy promontory develops pointing out towards the ocean. It is believed that this protrusion occurred after 1963 as it is not present in the lower quality 1963 headland photo. Post 1973 the promontory changes shape, perhaps through erosion on the

southern side. The variation in headland shape towards the south between 1973 and 1979 is likely to be a result of the fuzziness caused by sand and vegetation deposited on the rocks near the pier.

- 2.37 By 1985, the protrusion has been integrated into the headland with a much smoother headland boundary curving around to the north of the study site. This smoother shape remains constant till 1990, with strong overlap found between the 1985 and 1990 headlands. Between 1990 and the present, the middle of the headland accretes, expanding by around 10 metres.
 - 2.38 These data are interpreted further in chapter 7.
-

CHAPTER 3

RADIUM, ITS USE AND DISPOSAL

- 3.1 Radium-226 is a metallic radionuclide that emits alpha and gamma radiation and has a physical half-life of 1600 years. When it is taken into the body (by ingestion or inhalation), it is the alpha radiation that poses the greatest radiological hazard. The first product in its radioactive decay chain is the gas radon-222, which is itself a radionuclide (half-life of 3.8 days), also emitting alpha radiation (Figure 3.1). Other radionuclides in the decay chain emit alpha, beta and gamma radiation and include polonium-210 and lead-210; the last nuclide in the chain is stable lead-206. Radium-226 occurs naturally as part of the radioactive decay chain of uranium-238, and it can be extracted from uranium-bearing ores and purified by chemical means.




Figure 3.1 Decay chain of radium-226, showing radionuclide, type of radioactive decay and physical half-life

- 3.2 In the early 20th century, one of the major industrial uses of radium was in the production of luminescent paint. Fluorescence was achieved by combining radium-226 bromide and potassium sulphate solution with zinc sulphide crystals and glue was then added to produce paint. Whilst this was used in a number of products, its main application was associated with display dials and aircraft instruments (to make them visible in night flying). This process was known as 'luminising'. Another radionuclide, radium-228 (a beta-particle emitter) was used also, but was much less widespread. Due to its shorter physical half-life of 5.8 years, little of this radionuclide will now remain, even if it had been used on the site.

- 3.3 The majority of luminising carried out in the UK was before, during and for a short period after the Second World War, peaking during the war years. By the time the Radioactive Substances Act 1960 (RSA60) came into force on 1 April 1963, the luminising industry in the UK had dramatically diminished. Some commercial luminising was continued up to the late 1970s in support of military programmes and the watch industry. In addition, the military themselves, mainly the Royal Electrical and Mechanical Engineers (REME), continued to luminise and repair luminised items into the 1980s, but on a diminishing scale.
- 3.4 The diminishing quantity of luminous items represented a shift away from the perceived need of the military for dials to be luminous, accelerated by the introduction of VDUs and the replacement of direct luminous painting with gaseous tritium light devices from the late 1960s.
- 3.5 For aircraft instrument luminising, the typical radioactivity concentration of the paint was between 2.7 and 3.7 MBq per gramme (RSAC, 1958). Documents from Oak Ridge Associated Universities state that aircraft and ship instruments could contain 215 µg of radium per gramme of material to conform to British Admiralty standards, which would equate to 8 MBq per gramme. The former figure would refer to wet paint, while the latter would be the dry value and so the two are consistent.
- 3.6 Particularly following the Second World War, large numbers of military aircraft were scrapped and the instruments stripped out; it was common practice for these materials to be incinerated to reduce the volume of contaminated material and the resulting ash buried. One such location where both luminising and dismantling activities are known to have been carried out is at the airfield operated by the RNAS Donibristle and HMS Merlin. Wright (1990) notes that the MoD confirmed that some 800 aircraft were scrapped on the site in 1946. A witness statement obtained by SEPA (2013a) confirms that luminising activities on the site continued up to 1958. As noted above, the airfield closed in 1959.
- 3.7 The Radioactive Substances Advisory Committee (RSAC) noted in its 1958 report (paragraph 40) that *'the solid waste from normal operation is generally combustible in nature and it is fairly general practice to pour contaminated solvent, used for cleaning equipment, on paper and burn the whole under primitive conditions in the grounds. There is some local fall-out of ash; the petrol drum or old dust-bin used for burning the waste becomes fairly highly contaminated; and the ash remains for disposal. Non-combustible waste and the ash are generally buried in the grounds. Sometimes, and apparently usually at Services' workshops, all the solid waste is disposed of by burial'* (RSAC, 1958).
- 3.8 In terms of disposal, the RSAC report (paragraph 130) stated that *'although burial is more suitable for shorter-lived radionuclides, some with longer half-lives can be disposed of by burial provided it can be foreseen that the land will not be used for many years. We doubt whether it is practicable, except for a national site, to foresee possible future use sufficiently far ahead to justify disposal by burial of more than say a hundred microcuries of radionuclides of half-life longer than a year. Records of burials and burial sites should be kept and handed on to future users of the land.'* The 'hundred microcuries' referred to is equivalent to 3.7 MBq in modern SI units.
- 3.9 The hazards from radium presented by the residual ash relate to the potential ingestion or inhalation of, or skin contact with, contaminated material. A lesser hazard is that from external gamma radiation.
- 3.10 Radon-222 is the first decay product of radium-226. With its 3.82 day physical half-life, radon gas can diffuse through the ground before decay. If this occurs outdoors, it is diluted rapidly to

low concentrations. If buildings are erected over, or near to, a site of radium deposition, the radon gas can reach high concentrations indoors.

- 3.11 The average concentration of radium-226 in a home in the UK is about 20 Bq.m⁻³ ^a and the Advisory Group on Ionising Radiation concluded that around 1,100 radon-induced lung cancer deaths occur each year in the UK (AGIR, 2009). An HPA report stated that radon was the largest contributor to the average annual population radiation exposure, delivering some 1,300 µSv from a total of 2,700 µSv (Watson et al, 2005). The risk of developing lung cancer increases with increasing radon activity concentrations and is higher for smokers. The HPA (now PHE) has advised that indoor radon above an Action Level of 200 Bq.m⁻³ should be reduced.
- 3.12 For this reason, measurements of radon have been carried out by the HPA in buildings potentially affected. The precise values obtained remain confidential on grounds of personal privacy and were not reported to the Committee, which was assured that they were not significantly elevated.

^a <http://www.ukradon.org/article.php?key=radonaaffected>

CHAPTER 4

CONTAMINATION HISTORY

- 4.1 Following the discovery of radioactive contamination on the beach at Dalgety Bay during the routine baseline monitoring campaign by Babcock Engineering Services for the Rosyth Naval Base in June 1990, a limited survey was carried out which confirmed the presence of discrete sources of radium-226 on the beach. It was concluded that a more rigorous survey was necessary to determine the extent of the contamination.
- 4.2 In October 1990, a survey of the beach by the Directorate of Fisheries Research (Camplin, 1990) detected and recovered what was described as a particularly hot source from a depth of 15-20 cm. This source had an external dose rate of >28 mSv/h and its discovery resulted in a recommendation that access to the area should be restricted until a full radiological survey had been undertaken.
- 4.3 At the request of HM Industrial Pollution Inspectorate for Scotland (HMIPI), a survey was conducted during the latter half of November, 1990 by the then National Radiological Protection Board (NRPB)^b. This investigation indicated that all of the areas surveyed were contaminated with sources of radium-226. Consequently, it was recommended that the survey of the beach and foreshore be continued and extended to include the undeveloped land behind the beach and part of the housing estate close to the beach and foreshore.
- 4.4 With one exception (a fragment of painted metal found in a garden), all the radioactive contamination found consisted of radium-226 contained in, or associated with, clinker and ash material, and in general consisted of discrete sources. A total of 220 particles and 354 objects were recovered. The contamination was believed to have come from the disposal, by incineration, of dials from instruments painted with radium in the 1940s. The garden was remediated by the removal of some 2 tonnes of contaminated soil containing approximately 100 MBq of radium-226 and its replacement with new topsoil (Milton, 1991); a further 0.7 tonnes containing 4 MBq of radium-226 was removed following a re-survey. The total activity of the contaminated material removed during the survey has been estimated to be 135 MBq. This material was disposed of as radioactive waste by NRPB. Her Majesty's Industrial Pollution Inspectorate had already decided to continue to survey the beach and foreshore at 6 monthly intervals.
- 4.5 NRPB selected a representative sample of 24 private houses and direct radon measurements were made in living areas of these houses and surveys carried out of 17 gardens. The levels of radon were reported to be low. Contaminated material was found in two gardens.
- 4.6 The NRPB report concluded that the radiation dose to any individual as a result of ingestion of the material would be negligible; the material was deemed insoluble and as such would be eliminated from the body quickly (Milton, 1991).
- 4.7 The report was considered by COMARE in September 1991. The Committee was largely content with the actions and recommendations expressed in the report, but it asked for the following additional actions:
 - i) ISD should be asked to provide a report on the incidence of cancer in the local population,
 - ii) a reassessment should be made of the frequency of screening contaminated areas,

^b NRPB was subsequently incorporated into the Health Protection Agency (HPA), now renamed Public Health England (PHE)

iii) foundations of new buildings should be monitored for radioactivity.

- 4.8 On querying the possibility of other similar sites, the Committee was informed that MoD did not possess the data to give a comprehensive list and that it was probable that in the past disposal of luminised material would have been very widespread. A limited list of 26 sites (including Dalgety Bay) was supplied and the MoD confirmed that information on plants producing luminised material was well documented.
- 4.9 The epidemiology report from ISD was considered at the Committee's June 1992 meeting. It found that the only statistically significant differences observed were for pancreatic and non-melanoma skin cancers. Adjustments for socioeconomic factors using the Carstairs score (Carstairs and Morris, 1989) did not affect these findings. No overall excess of cancer was found in the Dalgety Bay area.
- 4.10 The Committee had no disagreement with the overall conclusions of the ISD report, and made the recommendation that the report should be published within the open scientific literature (see Black et al, 1994). It recommended also that it would be sensible to continue with the ongoing monitoring programme for the foreseeable future. Finally, it was recommended that Dalgety Bay should be re-evaluated by ISD in a few years' time, when the matter could be referred to COMARE.
- 4.11 In 1996, a survey of the surface gamma radiation dose-rate was carried out by the University of Aberdeen, covering '*representative sectors of the beach and foreshore, mud flats, the boat club area and the woods*' (Heaton et al., 1996). An average dose-rate of $0.088 \mu\text{Sv.h}^{-1}$ was found for the beach and foreshore, $0.081 \mu\text{Sv.h}^{-1}$ for the boat club and $0.066 \mu\text{Sv.h}^{-1}$ for the woods and mud flats. These results are not significantly different from the UK average for natural background gamma radiation (Watson et al, 2005). Therefore, even though several 'hot spots' with a reading generally 50% higher than the average were found, it was felt to be very unlikely that any deleterious effects would arise from enhanced exposure to external sources of gamma radiation at Dalgety Bay.
- 4.12 In June 2006, SEPA carried out a screening assessment of the area, which suggested that there was a realistic probability of harm occurring at Dalgety Bay due to the sources discovered. SEPA, supported by COMARE, recommended that further research be carried out into the risk from the sources and a desk exercise was undertaken by the MoD Defence Estates. The highest activity source recovered was 1 MBq of radium-226. There was no assessment of the distribution of contamination at that time, only an assessment of the risk of encountering the sources. As a result of this, signs were erected for the community promoting good hygiene, with recommendations to wash hands after visiting the beach. The Committee expressed concern over the timescale involved in the MoD work and any possible delay with remediation.
- 4.13 Subsequent surveys of the beach area were undertaken by various contractors for the MoD during which sources were removed from the beach to provide some protection to beach users. In 2007, the MoD employed Enviro Consulting to investigate the extent of radium-226 contaminated soils at Dalgety Bay, including the Sailing Club grounds, the gardens of 34 nearby residential properties and the coastal path. Radium-226 activity levels above background were identified on part of the coastal path, on Sailing Club land and in two gardens; the latter were remediated.
- 4.14 The Committee can only make recommendations and, until 2007, SEPA had no statutory powers to implement them. This position changed in 2007 with the introduction of the Radioactive Contaminated Land Regulations (Scotland), which placed a statutory duty on SEPA for sites such

as Dalgety Bay. The associated Statutory Guidance (Scottish Government, 2008) was introduced following these regulations.

- 4.15 In early 2008, following a report on the above work (Enviros, 2008), Enviro Consulting undertook supplementary investigations in the gardens of six properties located in the vicinity of the original MoD Salvage Yard, together with initial surveys on one property for which access was not previously available and two further properties at the request of SEPA based on the report's findings. Areas that were previously inaccessible at the sailing club were also assessed. Contamination was found in, and removed from, the gardens of six houses.
- 4.16 A COMARE working group visited the site in 2008 after completion of the remedial work and recommended that radon surveys of homes should be offered to those potentially affected by the new contamination finds. The beaches were examined and some concerns were raised, especially with regard to the integrity of the rock armour, as there was abundant evidence that quite large objects had moved from time to time, including the rock armour. While the Committee was content with the terrestrial work carried out by MoD, it recommended that more extensive surveys of the shore and inter-tidal areas should be carried out. The radon surveys were carried out and did not identify any raised levels of radioactivity.
- 4.17 SEPA and the MoD then agreed on a suitable monitoring periodicity. The 2008 SEPA survey detected 39 point sources (of which 16 were selected for leachate analysis) together with more diffuse radium contamination. The contamination appeared to be limited to an area between the old pipeline and the New Harbour.
- 4.18 Also in 2008, SEPA prioritised Dalgety Bay for assessment under the Radioactive Contaminated Land Regulations. In support of this, further monitoring, retrieval, sampling and analysis were carried out by SEPA. In response to their report, Defence Estates produced plans which could constitute an appropriate management plan, including an intensive monitoring and removal programme. The MoD met with SEPA, the HPA & the Food Standards Agency (FSA) to discuss plans for a full risk assessment.
- 4.19 Annual surveys continued during 2009 and 2010, but the Committee remained concerned about the inter-tidal area at Dalgety Bay. Defence Estates had not done any further characterisation and the annual monitoring programme had ceased. As a result, SEPA wrote to Defence Estates requesting them to undertake the work and, in the interim, took a sample of the point sources in order to fully characterise them and undertake work on the solubility of the radium-226 associated with the sources.
- 4.20 In 2011, SEPA undertook a physical examination of the headland area in an attempt to delineate the contaminated area and notified the Committee of an increased find rate of sources of higher activity than in previous years from particular locations. The Committee supported SEPA's decision to fence off a demarcated area, the establishment of a routine monthly monitoring programme in conjunction with the MoD, a request to the HPA for a further risk assessment and work to establish the solubility of the sources recovered.
- 4.21 In the year to October 2012, SEPA recovered over 800 sources from 36 one-day visits. It should be noted that the total area surveyed during these 36 visits did not cover the affected area in its entirety. The activity of sources recovered ranged from around 1 kBq to 76 MBq. Up to the end of 2012, several hundred additional sources were recovered by the MOD contractors, with around 100 sources being recovered each month over the 800 m coastline. Given that the population of sources had changed significantly, the gamma radiation dose-rate would have changed from that

previously measured also and so COMARE recommended that SEPA review this work and undertake further gamma monitoring as required.

- 4.22 It is the higher activity sources found over the last two years that could give rise to radiation doses of concern (Brown and Oatway, 2012; Dale, 2012). In addition, it was clear that there was ongoing repopulation (i.e. areas which had been surveyed and cleared of contamination exhibited further contamination within a short period of time).
- 4.23 There are several hypotheses as to the source of these re-occurring sources, as noted in Chapter 7. The situation appears to be a dynamic one, with the currently demarcated area being different from the focus of investigation just a year ago. Figure 4.1 shows the locations where the majority of the sources have been recovered.
- 4.24 Although a few larger items have been recovered, the vast majority of contaminated materials range in size from a few tens of millimetres to sub-millimetre and can be classed as artefacts associated with aircraft instruments and instrument panels through to clinker, indicating that the material has been incinerated. Chemical analysis shows that all sources have excess levels of zinc and sulphur, which can be attributed to radium paint.




Figure 4.1 Map of the Dalgety Bay site showing the distribution of radioactive items found by MoD and SEPA surveys in 2011 and 2012 (SEPA, 2013).

- 4.25 Table 4.1 shows the number of sources known to have been recovered from 1990 until the end of 2012.

Year	Number of sources removed
1990	226
1991	602
1992	108
1993	78
1994	34
1995	No monitoring data
1996	No monitoring data
1997	102
1998	11
1999	No monitoring data
2000	80
2001	No monitoring data
2002	93
2003	No monitoring data
2004	No monitoring data
2005	97
2006	37
2007	No monitoring data
2008	37
2009	104
2010	24
2011	530
2012	1369
Total	3532

Table 4.1 Beach monitoring finds (in 2009, 49 finds were a result of intrusive excavations).

- 4.26 In earlier reports, there appeared to be inconsistency in the source activities being reported by MoD contractors and SEPA. Once this had been resolved, COMARE's Contaminations Working Group suggested a standard for the detection limit to be used in the programme (20 kBq to a depth of 100 mm with 95% probability of detection), which was accepted by both parties and implemented in 2012.
- 4.27 The Contaminations Working Group considered whether the radium-226 activities reported by SEPA are correct. If these were overestimates, however, the values for source solubility reported below (paragraph 5.27) would increase, reaching values which would not be credible. Additionally, the solubility data were obtained from a UKAS accredited laboratory. It is unlikely, therefore, that there is any significant error.
- 4.28 The radium-226 activities of sources recovered since 2008 are shown in Table 4.2.

Source activity (kBq)	2008	2009	2010	2011	2012
No data	0	0	0	49	14
0-10	19	100	24	83	790
10-100	16	4	0	389	549
100-1000	2	0	0	4	15
>1000	0	0	0	5	1

Table 4.2: Source activity (using MoD contractor's conversion factors where appropriate)

Food & Environment Protection Act (FEPA) ban

- 4.29 In response to the 2011 source finds, the FSA carried out a screening assessment of the potential food chain dose based on the available data, and several results exceeded the 1 mSv annual allowable dose to members of the public, indicating a potential risk.
- 4.30 Responding to advice from FSA in Scotland, Scottish Ministers placed an Order under the Food & Environment Protection Act 1985 as a precautionary measure, restricting the removal of seafood from the area. Although there are no commercial fisheries in the area, individuals had been observed collecting shellfish, possibly for consumption. This was despite warning signs indicating that radioactive contamination had been found in the area and seafood should not be collected.
- 4.31 The Food Protection (Emergency Prohibitions) (Dalgety Bay) (Scotland) Order 2012 (SSI 2012 No. 135) (the 'FEPA Order') was placed on 9th May 2012 and approved by the Scottish Government Health & Sport Committee on 22 May 2012, making it an offence to remove fish & shellfish from the Dalgety Bay area.
- 4.32 Monthly sampling of cockles, mussels & winkles from Dalgety Bay was undertaken for a year from February 2012. The results for the first six months are shown in Table 4.3. No contaminated particles were detected in this exercise and this sampling has now been replaced by surface monitoring of the mussel bed areas.

SAMPLE	ANALYSIS RESULTS (Bq/kg)					
	Feb 2012	Mar 2012	Apr 2012	May 2012	Jun 2012	Jul 2012
Cockles	<10	n/a	n/a	n/a	n/a	n/a
Winkles (Site 1)	<10	<10	<10	<10	NS	NS
Winkles (Site 2)	<10	<10	<10	<10	<10	NS
Winkles (Site 3)	<10	<10	<10	<10	<10	NS
Winkles (Site 4)	<10	<10	<10	<10	<10	<10
Winkles (Site 5)	NS	NS	NS	NS	<10	NS
Mussels (Site 1)	<10	<10	<10	<10	NS	NS
Mussels (Site 2)	<10	<10	<10	<10	<10	NS
Mussels (Site 3)	<10	<10	<10	<10	<10	NS
Mussels (Site 4)	<10	<10	<10	<10	<10	<10

n/a – sample not available

NS – not sampled in order to maintain sustainability

Table 4.3 Dalgety Bay shellfish sampling.

- 4.33 The FSA in Scotland gave an undertaking to the Health & Sport Committee to review the FEPA Order in light of new evidence and/or effective remediation of the contamination. On 16 April 2013, a summary of monitoring undertaken and a restatement of this undertaking were provided to the Committee. This review is dependent on site remediation to be agreed between SEPA and MoD with whom FSA in Scotland will continue to liaise through the Dalgety Bay Particles Advisory Group.
-

CHAPTER 5

DOSIMETRY AND RISK

- 5.1 For radioactive material to pose any risk to the public there is a need for a pathway to exist between the source (particles) and a member of the public. These pathways are the route by which members of the public can become exposed to the source and thereby harm can occur. Whilst a radioactive source will present a hazard, it cannot pose a risk to the public without the existence of a pathway. The potential pathways of interest in the current situation are ingestion, inhalation, direct skin contact and external irradiation.
- 5.2 It is usual to consider that sources are encountered haphazardly or accidentally (i.e. on a chance basis), but some of the objects found in the area, such as luminised aircraft dials, would stand out from other items. The possibility of preferential selection, that someone would seek to investigate or collect such items, thus increasing the probability of encounter, must be taken into account when considering these pathways.

Habits survey

- 5.3 COMARE had recommended in 2008 that a habits survey should be undertaken. This was conducted at Dalgety Bay in October 2012 by the Centre for Environment, Fisheries and Aquaculture Science (Cefas) (Clyne et al, 2013) and involved the interviewing of people using the beach area. As the survey was undertaken in the autumn, it may have missed the higher occupancies and/or different habits of both tourists and the local population over the summer months.
- 5.4 The survey reported that there were many access points to the area and that walkers with and without dogs walked through part or all of the area. Many local walkers followed a regular route while others preferred to vary their route from day to day. Although most walkers reportedly stayed predominantly on the paths and grass areas above the shore, approximately 60% of the dog walkers and 30% of the walkers without dogs regularly visited one or more of the beaches as part of their walk.
- 5.5 Several families spent time on the small sand beach to the west of New Harbour, engaged in activities such as playing and building sandcastles or rock pooling in the rocks nearby. They reported that they also visited this beach in the summer time, when they took picnics and paddled or swam in the sea. A few families also played and paddled on the small sand beach at New Harbour and one family was identified who spent a small amount of time playing on the shore at the Slipways. The families were noted to include very young children.
- 5.6 A few individuals who were walking, dog walking or playing, picked up shells or stones from the beaches at Ross Plantation, the Slipways, New Harbour and the West End. Some people took the stones and shells home with them. No one was seen to be deliberately searching for old aircraft artefacts on the shore at the time of the survey, although this has been reported to occur (McPhail, 2011b).
- 5.7 A small number of runners and joggers were recorded and they took similar routes through the survey area to the walkers. A few cyclists passed through all or part of the survey area and they stayed on the paths and grass areas above the shore. Horse riders were observed on the Fife Coastal Path, but were not interviewed. Four bird watchers were interviewed. They mainly used the beach at Ross Plantation since it provides a good vantage point across the mudflats of the bay,

which attract many birds. The bird watchers also spent time on the grass areas and paths above the shore at the Headland and in other zones.

- 5.8 Several people commented that the sand beach at West End could be very busy with families having days out on the beach in the summer, and that the Fife Coastal Path attracted more walkers in the summer months.
- 5.9 Boating activities centred on the sailing club. The club caters for dinghy sailors, keelboat sailors and users of powerboats, mainly rigid inflatable boats (RIBs). Adults and children sail at the club. The main sailing season was found to be between April and October although a minority of sailors continued through to December. One individual was identified who used a powerboat throughout the year. The club ran regular racing programmes for dinghies and keelboats and sail training courses. During the summer they held a Youth Sailing Week event to encourage youngsters to sail. Most dinghy sailing took place in the waters close to the club. The keelboats and powerboats also sailed locally, but also went further afield in the Firth of Forth, and some keelboats went cruising to areas outside the Firth of Forth. In 2012, a large number of the Club's regattas and events were cancelled; as a result the numbers of people using the slipway areas was significantly lower in 2012 than it had been in previous years. Local people have reported that the regattas had been very popular, often resulting in crowded beach areas.
- 5.10 Boat maintenance was identified as taking place on the land close to the shore in the Boat Area zone, on the fringe of the upper shore near the main slipway and on the beach at New Harbour. Boat maintenance included work carried out on the club raft as well as on boats. Keelboats were kept moored offshore from the sailing club and most boat owners lifted their moorings once or twice a year to check the condition of the chains and shackles.
- 5.11 Several people reported that they had seen one or two people digging for angling bait in the area of mud, sand and stones on the lower shore of the West End zone. This activity was reported to take place only around the time of spring tides (i.e. fortnightly) when more of the shore was exposed at low tide. This activity was observed during the physical investigations at Dalgety Bay. It was reported also that gangs of commercial winkle pickers had occasionally been seen on the shore, both within the survey area and also just outside the survey area on the shore near St. Bridget's Kirk, further east in Dalgety Bay. However, interviewees were uncertain how long ago they had last seen this activity taking place.
- 5.12 It was reported that one individual spent a number of nights sleeping on the beach at Dalgety Bay, and although this was not observed during the habits survey itself, the individual is known to the council as this activity occurs periodically. In 2011, SEPA also received a call regarding the presence of a scout group using the now demarcated area for recreational purposes.
- 5.13 In 2010, following recovery of sources from the slipway area, SEPA requested that the local pre-school playgroup cease using the location and move to the New Harbour. Following the identification of radioactive contamination at the latter location also in 2012, the playgroup was advised to move to an area further to the west where no contamination has been detected.
- 5.14 Having established the various groups who might encounter sources, it is necessary to establish the consequences of the routes by which the public could be exposed.

Inhalation

- 5.15 For inhalation to be a hazard, the source would have to be a very small particle. The radiation dose from a given source activity increases with the depth in the respiratory tract at which deposition occurs. This depth increases with decreasing particle size.

- 5.16 Particles of more than a few tens of μm have a very low probability of reaching the airways and alveolar region of the lungs – if inhaled, they are likely to be trapped in the nose and extrathoracic airways.
- 5.17 It has been suggested (Breslin, 2011) that particles of radium of a size which could be inhaled would be present at Dalgety Bay.
- 5.18 In 2012, the HPA reported that “*for a particle containing 100 Bq of radium-226 and a diameter of 100 μm the committed effective dose would be around 0.1 mSv*”. Such a dose would be a low hazard if encountered in isolation (Smith, 2010).
- 5.19 Wilkins (2008) calculated that if a particle containing 1 kBq of radium-226 was sufficiently small to be deposited in the alveolar regions of the lung, the committed effective dose would be 1-25 mSv for an adult or 50-150 mSv for a 1 year-old child. The doses would be higher for bronchiolar deposition.
- 5.20 Heaton et al (1996) reported, however, that less than 1% of a sediment sample from the beach was composed of particles less than 45 μm and stated that “*there would be a low risk of members of the public inhaling fine material*”.
- 5.21 While all of the sources recovered in recent months have been of a size greater than 100 μm , several have been shown to be friable. This raises the possibility that, over time, larger sources containing high levels of radium-226 could break down into a large number of lower activity sources which would not be detectable by the current routine monitoring equipment, but would be capable of being inhaled.
- 5.22 In order to exclude the possibility of such a mechanism, SEPA examined a number of grab samples of surface material which were assayed on a more sensitive detector. No evidence of the presence of low activity sources was found.
- 5.23 On the basis of the above information, the Committee is of the view that there is no reason to believe at present that the inhalation pathway is significant in terms of public health at Dalgety Bay.

Ingestion

- 5.24 The most important variable in calculating the radiation dose from ingestion is the solubility of the material, since this determines the fraction of radium-226 transferred into the bloodstream and thence to body organs.
- 5.25 With support from COMARE (see paragraph 4.19 above), SEPA contracted the University of Stirling to carry out a study of the solubility of a representative sample of sources. Sources were digested in solutions mimicking stomach and small intestine (Tyler et al, 2013), following the protocol established for Sellafield and Dounreay (Harrison et al, 2005).
- 5.26 A sample of sixty sources were selected, on the basis of their activity and dimensions, and subjected to digestion in simulated stomach and lower intestine solutions. The study demonstrated that significantly more radium-226 and lead-210, driven by polonium solubility, was dissolved from sources in ‘stomach’ solutions compared with ‘lower intestine’ solutions. The combined ‘gut’ solubility for radium-226 and apparent lead-210 varied from less than 1% to up to 35%. The difference could be due to the effects of burning at differing temperatures affecting the chemical form of the radium and thus the solubility. For example, burning may result in the production of radium oxide which would be highly soluble in the gastrointestinal tract.

- 5.27 The solubility results were not normally distributed and so a log transformation was applied. Following Gilbert (1987) this corrected estimate for mean and standard deviation gives values of 7.6% and 2.5% respectively, with a 95th percentile value of 20.2%. If a cautionary approach were to be adopted, a reasonable level of caution would be to use the 95th percentile for the assessment of hazard.
- 5.28 In 2013, the Committee's Contaminations Working Group considered the possibility that the *in vitro* measurements might overestimate the mammalian *in vivo* value. It was noted that, due to the apparent differences in chemical form of the sources found, it would be necessary to test large numbers of animals to resolve this issue. Use of the relatively low mean value from the distribution already makes some allowance for this uncertainty.
- 5.29 Radiation doses were derived using standard ICRP dose coefficients (ICRP, 1996). Assuming that the radium-226 daughters are in equilibrium, the contributions from radium-226, bismuth-214, lead-214, lead-210, bismuth-210 and polonium-210 are included in the calculation and bismuth-210 and polonium-210 are assumed to be in equilibrium with lead-210. This is an important assumption as 66 % of the estimated dose for an infant less than 1 year of age is dominated by polonium-210 versus 55 % for an adult. This assumption has been validated by results from a recent intercomparison exercise by the Terrestrial Environment Laboratory of the International Atomic Energy Agency (IAEA, 2012).
- 5.30 Tables 5.1 and 5.2 show the results for mean and 95th percentile solubilities respectively. Table 5.1 shows that sources which have the mean solubility would require activities of around 100 kBq of radium-226 to deliver a committed effective dose in excess of 100 mSv to a one year old child (age range 1 to 2 years). A little over 10% of the sources tested dissolved sufficient radioactivity to result in 100 mSv committed effective dose to an infant. If the 95th percentile value is used, sources of greater than 40 kBq would result in a dose of greater than 100 mSv to a one year old child and for an adult this would be around 250 kBq.

		Committed effective dose (mSv)					
Original Activity (Bq)	Activity in sol. (Bq)	3 months	1 year	5 years	10 years	15 years	Adult
1,000	76	3	1	0.55	0.40	0.38	0.16
10,000	760	30	10	5.5	4.0	3.8	1.6
100,000	7,600	300	100	55	40	38	16
1,000,000	76,000	3000	1000	550	400	380	164

Table 5.1 Doses arising from ingesting a source of given activity with mean solubility of 7.6%.

Original Activity (Bq)	Activity in sol. (Bq)	Committed effective dose (mSv)					
		3 months	1 year	5 years	10 years	15 years	Adult
1000	202	7.9	2.7	1.5	1.1	1.0	0.44
10000	2,020	79	27	15	11	10	4.4
100000	20,200	790	270	150	110	100	44
1000000	202,000	7900	2700	1500	1100	1000	440

Table 5.2 Doses arising from ingesting a source with solubility at the 95% of the distribution (20.2%).

- 5.31 Some of this information together with further unpublished information was used in an HPA report on the hazards posed (Brown & Oatway, 2012). In that report, the HPA assumed that the maximum activity which could be ingested was 100 kBq radium-226, with a physical diameter of 1 mm. Litovitz et al (2010) reported, however, that batteries of up to 20 mm in size have been implicated in many of the complications from button battery ingestions by children less than 4 years of age. Thus it is possible that sources of a few mm diameter would be ingestible and these could have an activity of around 1 MBq. Indeed, sources have been recovered at Dalgety Bay which have multi-megabecquerel activities and physical sizes less than 20 mm.
- 5.32 In the extreme case of ingestion of the highest activity source recovered, the 2012 HPA report stated that the possibility of acute organ damage or bone marrow failure cannot be excluded, but the majority of recovered finds have activities much less than this value (Brown & Oatway, 2012). The HPA report then concludes, however, that the committed effective dose from ingestion of a 100kBq source could be around 330 mSv to a young (1-2 year old) child, similar to values derived above. For an ingestible 1 MBq source this would suggest a dose of a few Sv for a young child, which could lead to the acute consequences outlined above.
- 5.33 As indicated above, only a few sources of 1MBq or greater have been found to date and thus the current possibility of this occurring would be remote using present beach find data especially as, at present, the known contaminated area is under monthly surveillance. Nevertheless, even the potential radiation doses from sources of 100kBq are significant.
- 5.34 Samartzis et al. (2011) investigated bone cancer incidence among the Japanese atomic bomb survivors, who were exposed largely to an external source of gamma radiation. They found an excess incidence of bone cancers among survivors followed to 2001, but only at bone marrow doses estimated to be greater than 0.85Gy; above this value, the dose-response relationship was linear with an excess relative risk of 7.5 per Gy (95% confidence interval: 1.34, 23.14). In its 2006 report, UNSCEAR (2008) modelled the radiation-induced excess risk of bone cancer incidence among the atomic bomb survivors followed to 1998 and found that the excess risk was best described by a quadratic dose-response – the excess absolute risk (EAR) was best described by $9.33 \times 10^{-6} \times D^2$ cases per 10^4 person.year-Sv, where D is the equivalent dose to the skeleton. Follow-up of solid cancer incidence among the atomic bomb survivors, however, only commenced in 1958, 12.5 years after the bombings, so these investigations would have missed any early radiation-induced bone cancers. The work of Nekolla et al. (2000), upon which the United States Environmental Protection Agency (EPA) bone cancer incidence risk model is largely based, suggests that such early bone cancers would have occurred. The Nekolla et al.

(2000) study is of patients injected with radium-224, which emits alpha-particles that irradiate the bone surfaces (as will occur also when radium-226 is deposited on bone surfaces) and so leads the Committee to base its bone cancer risk estimates upon the EPA bone cancer incidence risk model (EPA, 2011; Pawel and Puskin, 2012). The nominal bone cancer incidence risk estimate used by the ICRP (2007) of 6.5×10^{-4} cases per person-Sv is also based on studies of the radium-224-injected patients, as is the bone cancer mortality risk estimate of 5×10^{-4} deaths per person-Sv derived by the former National Radiological Protection Board (NRPB 1993).

5.35 The EPA bone cancer risk model takes the form

$$EAR = \alpha D g(e) h(t) \text{ excess cases per } 10^4 \text{ person-Gy}$$

where D is the bone surface dose of alpha-particle radiation in gray

$$\alpha = 1.7 \times 10^{-3} \text{ Gy}^{-1}$$

$$g(e) = \exp[-0.0532(e-30)]$$

$$h(t) = (2\pi\sigma^2)^{-1/2} \times \exp[-(\ln(t)-\ln(t_0))^2/2\sigma^2] \times 1/t$$

e is age-at-exposure in years

t is time-since-exposure in years

$$\sigma = 0.612$$

$$t_0 = 12.72 \text{ years.}$$

Thus the EPA model describes an excess absolute risk of bone cancer incidence that is linear with the bone surface dose of alpha-particle radiation (and without a threshold dose), and is modified by the age-at-exposure (greater at younger ages at exposure) and by time-since-exposure (excess risk is expressed as a “wave” with time-since-exposure, peaking at nine years since exposure). The EPA (2011) suggests that this model may be conservative (in that it somewhat overestimates the risk), a position also adopted by the ICRP (2007) for its nominal risk coefficient for bone cancer.

- 5.36 At the request of the Committee, Public Health England provided estimates of the absorbed doses to various tissues received in each year following the ingestion of 1 Bq of radium-226 at ages 3 months and 1, 5, 10, 15 years, and by an adult using the ICRP solubility assumptions. The doses are dominated by the alpha-particle dose, and the dose to the bone surface is substantially greater than that to other tissues (e.g. the bone surface dose is some 40 times greater than the liver dose). The bone surface doses are highest for an infant aged 3 months, and decrease with time-since-exposure (as radium-226 becomes buried in the bone volume). For an infant ingesting 100 kBq of radium-226, the bone surface dose from alpha-particles received to one year of age is 0.66 Gy, and the lifetime dose is 0.84 Gy. Applying the EPA bone cancer risk model to these bone surface doses received following intake in infancy gives a lifetime risk of bone cancer incidence of 0.86%. The ICRP nominal risk coefficient also gives a lifetime risk of around 1%, while the UNSCEAR bone cancer risk model gives a lifetime risk less than that given by the EPA model by a factor of 3½. Since the EPA bone cancer risk model has a linear, no-threshold dose-response, intakes of other activities of radium-226 will result in risks that are directly proportional to that arising from an intake of 100 kBq. The lifetime risk of bone cancer incidence following ingestion of radium-226 at older ages will be less than that for an infant, substantially so for adults.

Skin contact

- 5.37 The main deterministic effect of importance is ulceration, for which cells at greater depths in the dermis are implicated. For adults, the relevant depth is taken to be 70 μm , at which there is no contribution from alpha radiation.
- 5.38 Using a standard computer model, Charles and Gow (2010) estimated that the total skin dose rate (at a depth of 70 μm over an area of 1 cm^2) from a radium-226 point source activity of 1 MBq (in equilibrium with all progeny) is approximately 5.5 Gy.h^{-1} . A theoretical value of 3.8 Gy.h^{-1} has been obtained by PHE (Tanner, 2013). Taken together, it would be reasonable to assume a maximum dose rate of 5 Gy.h^{-1} . For the point sources recovered this is likely to be an over-estimate of dose by about 10% since progeny in the chain following radon-222 will be in deficit relative to the equilibrium distribution. The main contributors to the dose are lead-214, bismuth-214 and bismuth-210; the last of these will be in deficit due to the 22 year physical half-life of its precursor, lead-210. These theoretical dose rates will be affected also by source shape and self-absorption characteristics.
- 5.39 Using radiochromic dye film measurements (Charles and Gow, 2010), experimental estimates of dose were obtained from ten sources. Results from the two highest activity sources are likely to have been the most reliable. One showed the dose rate at 70 μm depth to be 1 Gy.h^{-1} per MBq, whilst the other gave a value of 2 Gy.h^{-1} per MBq. The latter is consistent with a report from the MoD (Brown, 2008), which estimated the skin contact dose to be 2.7 Gy.h^{-1} per MBq.
- 5.40 The 2012 HPA report recommended the use of a value of 1 Gy.h^{-1} from a 1 MBq source on the basis of theoretical calculation (Brown & Oatway, 2012).
- 5.41 Experimental studies of dose rates from the 76 MBq source recovered at Dalgety Bay in 2011 have been carried out recently (Burgess, 2013). These demonstrated that the measured dose rates around the object varied by at least three orders of magnitude. This would suggest a localised source of paint within a matrix of either inert or much lower level radium contamination substrate. The maximum dose rate equivalent reported at 70 μm depth was 1,380 mSv.hr^{-1} averaged over the 4.5 mm diameter thermoluminescent dosimeter. This would suggest that dose rates of around 1-2 Gy.h^{-1} are possible from specific locations in even such a large source. As the measurements were located randomly, it is possible that higher doses could occur if this location were not the highest concentration, but it is unlikely to be much greater since 22 measurements were made at locations across the surface of the source and the next highest reading was 286 mSv.h^{-1} , obtained within a few mm of the highest value. However, if the activity being measured is subsurface, then physical breakdown of the source could result in significantly greater dose rates.
- 5.42 The above values are for adult skin, but in young children reference skins are thinner (45 μm) than in adults (70 μm) (ICRP, 1975; ICRP, 2002). Theoretical calculations by Charles and Gow (2010) indicated that, for skin depths of less than 70 μm , the predicted depth dose curves showed a sharp increase due to alpha particle dose. For depths between 40-50 μm , the calculated alpha absorbed dose was more than an order of magnitude greater than the beta/gamma absorbed dose. Their experimental results, however, did not seem to accord with the prediction.
- 5.43 Overall, the available data indicate that it would be reasonable to assume that physically small sources could give rise to skin doses of 2 Gy.h^{-1} per MBq, while physically larger sources of very high activity could yield skin doses of several Gy.h^{-1} . Given possible residence times of more than an hour, high skin doses (in excess of the threshold of 2 Gy for deterministic effects) from several of the sources that have been recovered are a definite possibility.

External irradiation

- 5.44 The external irradiation hazard is due to the gamma radiation emitted from the sources and could arise from passing through the affected area or from lying on the ground near to a source.
- 5.45 As a result of the COMARE recommendation (see paragraph 4.20), SEPA undertook a gamma survey in order to provide an estimate of the radiation dose that could be received by people walking across the beach. Gamma dose rates were measured at 263 locations across the affected area. The maximum dose rate was found to be 73 nSv.h^{-1} (mean 27 and standard deviation 9.7 nSv.h^{-1}). Even if a person remained at that location for an entire year they would receive a dose of only 0.6 mSv; a more realistic 1h per day spent moving through the area would give rise to a dose of 10 μSv over a year using the mean radiation level measured. By comparison, the UK average background dose rate is some 2.7 mSv with the natural contribution to the total being around 2.2 mSv (Watson, 2005).
- 5.46 Assuming a direct relationship between activity and dose rate (i.e. neglecting geometrical effects), sources that contain MBq activities could deliver external doses of $>7.5 \text{ } \mu\text{Sv.h}^{-1}$ if a person came within 0.5 m of the source. However, the effects of shielding from burial of the source would reduce this significantly. If the maximum residence time for a person on the beach is around 8 hours (due to tides) and a person were to remain static for this period and within 0.5 m of a 70 MBq source on the surface of the beach, they could receive a dose of 1 mSv.
- 5.47 Given this calculation, the risk from external irradiation is not of major concern.

Groundwater

- 5.48 There is a theoretical possibility of radium leaching into groundwater from the contaminated material. The MoD contractor has taken five samples from the area and reported a result of 65 counts per minute (AMEC, 2013a). The import of these data is unclear since there were no values given for a control or for the minimum detectable level. Further information has been sought from the MoD.

CHAPTER 6

CANCER EPIDEMIOLOGY

- 6.1 Exposure to moderate and high levels of ionising radiation is known to cause most types of cancer (IARC, 2012; ICRP 2007). Not all types of cancer are equally sensitive to induction by radiation, however, and Boice (2006) has categorised cancers into four groupings:
- Cancers frequently associated with radiation with authoritative risk estimates
 - Cancers occasionally associated with radiation with robust risk estimates
 - Cancers rarely associated with radiation with uncertain risk estimates
 - Cancers never or sporadically associated with radiation with no risk estimates

By international consensus, the risk estimates obtained from the epidemiological study of groups exposed to moderate and high doses of ionising radiation (such as the Japanese atomic-bomb survivors) are assumed, for the purposes of radiological protection, to extend to low doses (ICRP 2007).

- 6.2 The risk of cancer posed by exposure to ionising radiation depends on the types of cell that are irradiated, the level of exposure and the sensitivity of the cells to radiation-induced cancer. Penetrating gamma radiation essentially irradiates all tissues, but short-range radiations, such as alpha-particle radiation, only irradiate those cells close to the site of deposition of the radioactive material. This is especially so for alpha-particle-emitting material deposited within the body. Therefore, if an alpha-emitter is distributed heterogeneously within the body at a high level, a characteristic pattern of radiation-induced cancer sites would be expected, reflecting the particular sensitive cells that have been irradiated by the alpha-particles.

Effects of radium

- 6.3 Radium is an alkaline earth metal and when taken into the body, it behaves biochemically like calcium, so that it is deposited predominantly in bone, initially with a brief residence on bone surfaces and then in the bone volume. Consequently, any health effects arising from the intake of radioisotopes of radium would be anticipated to be related to its primary site of deposition within the body, bone surfaces, and then bone volume.
- 6.4 Workers in the past, usually young women, who applied radium-based luminous paint to watch and instrument dials inadvertently ingested radium, mainly through the action of “tipping” by which they licked the tip of the paintbrush to obtain a fine point. Women in the luminising industry in the USA prior to 1930 were particularly heavily exposed to radium in this way and it has been known for many years that they experienced a substantial excess risk of bone and head cancers as a result of their ingestion of radium (e.g. Jones and Day, 1945).
- 6.5 Stebbings et al (1984) examined data on cancer mortality amongst female radium workers in the USA. In addition to the long-established excess mortality from bone and head cancers, they examined mortality from other cancers and found no clear evidence of an association between these other cancers and exposure to radium. In particular they noted that *‘liver, pancreatic, cervical, and uterine cancers were clearly unrelated to radium exposure’*.
- 6.6 In a comprehensive review of the evidence from studies of the US dial luminisers, Fry (1998) noted the large excess risk of bone cancers resulting from radium deposited in the bone. Remarkably, there was also a large excess risk of cancers of the paranasal sinuses and mastoid air cells in the head. This excess was attributed to the decay of radium-226 in the bones of the head

into the noble gas, radon-222, that was released into the airways of the head, leading to their irradiation by radon-222 and its radioactive decay products. Fry noted that although increased risks were observed for some other health outcomes, such as breast cancer, *'no significant or unequivocal relationships with prior exposure to radium have been found'*. Of some interest is the observation that all of the bone cancers occurred in workers with bone surface doses greater than 5 Gy (of predominantly alpha particle radiation), i.e. very high localised doses.

- 6.7 During the 1940s and 1950s, patients in Germany suffering from bone disease - primarily bone tuberculosis or ankylosing spondylitis, but including some patients with other medical conditions (e.g. polyarthritis) - were injected with radium-224 as a treatment. Radium-224 has a relatively short half-life of 3.6 days, decaying through the emission of an alpha-particle to produce another isotope of radon, radon-220. Radium-224 occurs naturally on Earth as a component of the radioactive decay chain of thorium-232. Owing to its short half-life, radium-224 decays mainly on the surfaces of bones where it is deposited, rather than in the bone volume.
- 6.8 Nekolla et al (2010) have reviewed the findings of the study of patients injected with radium-224. They found the anticipated large excess risk of bone cancers, but also reported excesses of a number of other cancers, such as thyroid, liver and breast. The standardised incidence ratio (SIR) for liver cancer was 4.2 (10 cases observed against 2.4 expected). Considerable care has to be exercised when assessing the results of studies of patients who have been treated with radiation as therapy for a disease because of concerns that the disease itself may have increased the risk of future diseases such as cancer. In this respect, it is of interest that the excess of cancers other than bone cancer is confined to those patients with bone tuberculosis, suggesting that this group of patients may be at an increased risk of these other cancers for reasons other than the radium-224 injections. Further, Nekolla et al (2010) note that pre-existing liver disease (present in 4 of the 10 patients) is likely to be a factor in the reported excess of liver cancers. Since radium-224 was administered as a therapy, the doses delivered to bone surfaces, at several Gy, were high.
- 6.9 In addition to being a different radioisotope of radium, the radium-224 administered to the patients was in a different chemical form from that used by the dial painters, being designed to be soluble in the bloodstream and deposited rapidly on bone surfaces. Doses to tissues other than bone surfaces were also high (e.g. the average dose to the liver was several Sv (Nekolla et al, 2010)) so that a radiation-induced excess of cancers other than bone cancer is plausible. Further, phenylbutazone was administered to patients to relieve the pain associated with bone disease, a drug that may be related to some cancers, such as acute leukaemia.

Overview

- 6.10 It is clear that intake of radium leads to an excess risk of bone cancers, but that this occurs after high alpha-particle doses to bone surfaces. Large intakes of radium-226 also lead to an excess risk of rare head cancers due to the production of radon-222 from the decay of radium-226 in the bones of the head.
- 6.11 The risk of other cancers from exposure to radium-226 is not well established, and for liver cancer in particular the small number of excess cases reported from the patients injected with radium-224 could well be due to other causes. An excess of liver cancer in the dial luminisers with high intakes of radium-226 has not been found.
- 6.12 No papers have been found in the peer-reviewed literature that reliably link the risk of non-Hodgkin lymphoma (NHL) to radium exposure.

ISD reports

- 6.13 Following receipt of the 1991 NRPB report (Milton, 1991), COMARE first commissioned a report from ISD on cancers in the Dalgety Bay area in 1992, which covered the period 1975-1990 and was subsequently published in 1994 (Black et al, 1994). The ISD study was extended in 2005 to cover 1975-2002. The most recent report for COMARE covered 2000-2009. The Committee had requested consideration of all sites of cancer in this report, which led to a large number of statistical tests comparing expected and observed numbers. For this reason, a higher than usual threshold was used to determine statistical significance, thus reducing the likelihood of finding a nominally positive statistical association by chance alone and significance tests were conducted at the 1% level. Also, since half of the postcodes involved are in the least deprived 10% of the Scottish population and all are within the least deprived 40%, the two least deprived deciles of the Scottish population were used to calculate the expected cancer rates. This partially corrects for known socioeconomic effects on cancer incidence rates (COMARE, 2006).
- 6.14 Data from the 1975-1990, 1975-2002 and 2000-2009 reports are shown in Table 6.1. It will be seen immediately that, for most sites of cancer, the average annual expected numbers of cases are substantially higher in the later period. This could merely reflect the fact that the population was predominantly younger in the last quarter of the 20th century, consisting as it did of incoming workers with young families (Black et al, 1994).
- 6.15 A statistically significant ($p<0.01$) excess of liver cancer incidence was present in 2000-2009, but not during 1975-2002, when there was a (statistically non-significant) deficit of cases. Liver cancer is not among the ten most common cancers for either men or women in Scotland, but the annual number of cases rose from 151 to 258 between 2000 and 2009 for men, although remaining at 112 per year for women over this period.
- 6.16 There was also a statistically significant ($p<0.01$) excess of cases of NHL during 2000-2009, but not in 1975-2002. The excess of cases of non-melanoma skin cancer (NMSC) during 1975-2002 is statistically significant ($p<0.01$), but the observed number of cases of NMSC is similar to the expected number during 2000-2009. The excess of cases of pancreatic cancer during 1975-1990 approaches statistical significance at the $p=0.01$ level, but over the period 1991-2009 the observed number of cases is about the same as the expected number.
- 6.17 These nominally statistically significant excesses of particular sites of cancer must be set against what is known about their causes and how they might be related to exposure to radium-226 at Dalgety Bay. The absence of any case of bone cancer during either period is notable, since bone cancer (and rare head cancers) is the only type of cancer firmly associated with exposure to radium-226; the pattern of cancers shown in Table 6.1 must be viewed in this light.
- 6.18 The excess of cases of Hodgkin's lymphoma during 2000-2009, which approaches statistical significance at the $p=0.01$ level, may suggest that there is a common cause for Hodgkin's lymphoma and NHL. Hodgkin's lymphoma is considered to have a very low sensitivity to induction by radiation, if it can be induced by radiation at all (Boice, 2006).

		1975-1990 ^c		1975-2002			2000-2009		
		Obs	O/E	Obs	Exp	O/E	Obs	Exp	O/E
Frequently associated with radiation	Myeloid Leukaemia (C92)						6	4.9	1.23
	Leukaemia (C91-C95)	5	1.02	15	15.32	0.98			
	Thyroid (C73)	3	2.4	6	3.76	1.6	3	2.9	1.03
	Female breast (C50)	31	1.19	92	84.6	1.09	61	73.8	0.83
Occasionally associated with radiation	Lung (C34) (C33-C34)	25	0.65	74	113	0.65	41	43.9	0.93
	Stomach (C16)	4	0.4	10	27.04	0.37	6	10	0.6
	Colon (C18)	15	1.06	44	46.61	0.94	34	38.9	0.87
	Oesophagus (C15)			7	15.3	0.46	6	10.4	0.58
	Bladder (C67)	8	0.86	22	26.97	0.82	9	10.4	0.86
	Ovary (C56)			12	14.87	0.81	9	10.7	0.84
	Brain and nervous system (C71-C72) (C70-72)			11	11.46	0.96	6	7.7	0.78
	Liver (C22)	0	0	2	4.55	0.44	10	4	2.47
Rarely associated with radiation	Kidney (C64) (C64-C65)	3	0.78	7	12.01	0.58	8	10.2	0.78
	Salivary glands (C07-C08)			2	1.57	1.27	2	0.5	3.65
	Non-Hodgkin lymphoma (C82-C85)	4	0.75	18	17.33	1.04	27	15.8	1.71
	Myeloma (C90) (C88+C90)	2	1.08	8	6.51	1.23	5	6.1	0.82
	Malignant melanoma (C43)			21	15.94	1.32	20	22.7	0.88
	NMSC (C44)	36	1.5	129	101.6	1.27	169	163.2	1.04
	Rectum (C19-C20)	10	1.36	32	23.78	1.35	18	19.9	0.9
	Uterus (C54-C55) (C54)			16	9.06	1.77	12	9.7	1.23
	Bone (C40-41)	0	0	0	1.59	0	0	0.4	0
	Connective tissues (C47+C49)			3	3.43	0.88	2	2.9	0.69
Never or sporadically associated with radiation	Chronic lymphocytic leukaemia (C91.1)						7	4.6	1.51
	Pancreas (C25)	11	2.28	18	14.42	1.25	12	9.3	1.28
	Hodgkin's disease (C81)	1	0.45				7	2.6	2.68
	Prostate (C61)	5	0.68				61	52.4	1.17
	Testis (C62)						3	4.2	0.71
	Cervix (C53)						3	3.6	0.83
	Selected childhood cancers (0-14 age range) ^d						0	0.4	0
Childhood leukaemia				4	2.12	1.89			
Other sites		48	0.93						
Total figures for cancers shown		Obs 211	Exp 214.2	Obs 553	Exp 572.8		Obs 547	Exp 546.1	

Table 6.1: Cancer incidence figures for 1975-1990, 1975-2002 and 2000-2009 in the Dalgety Bay area. Figures in bold are statistically significant increases in the observed number of cases over those expected, using a one-sided test at the 0.01 significance level.

^c Reference – Black et al (1994).

^d Sites for which radiation-induced cancers have not yet been reported or confirmed: retinoblastoma, Wilms' tumour, neuroblastoma, and others of embryonic origin (Boice, 2006)

- 6.19 The excess of non-melanoma skin cancer (NMSC) during 1975-2002 is comparable to that seen during the same period for melanoma skin cancer, a cancer with a very low sensitivity to induction by radiation (Boice, 2006). The excess incidence of melanoma and non-melanoma skin cancers may indicate a link to UV exposure, which is an established cause of skin cancer of both forms. The age distribution of the population at this early period is consistent with uptake of package holidays which could have led to excessive exposure to sunshine, although the incidence of both types of skin cancer returned to expected levels during 2000-2009.
- 6.20 The generally low ratios of observed to expected cases presented in Table 6.1 for those sites of cancer most strongly associated with radiation exposure by previous epidemiological studies (e.g. lung and oesophagus) are notable. The low rate of lung cancer argues against any significant exposure through inhalation of airborne sources.

Postcode analysis

- 6.21 Since the liver cancer and NHL incidence data from ISD demonstrate statistically significant increases in incidence, the Committee undertook a further analysis using areal counts in each of the 13 data zones surrounding the site, supplied by ISD. These zones are bounded by the Firth of Forth and A921 on the south and north, by Letham Hill to the west and the fields above Braefoot Point to the east.
- 6.22 Having first confirmed through a test allowing for the large number of cancer sites examined that the excess in incidence of both diseases was unlikely to be due solely to this multiplicity, the data were subjected to the 'Linear Risk Score' distance test, found previously (COMARE, 2005) to be well suited to datasets involving small numbers. In effect, the procedure tests whether the cases observed are distributed closer to a specific point – in this case the headland, where the contamination has been confirmed – than would be expected by chance alone.
- 6.23 There are two forms of this test, conditional and unconditional. The conditional test is used to assess whether there is a significant association between incidence and distance from the hypothetical source, taking no account of the overall incidence rate, while the unconditional test integrates the spatial evidence with the overall incidence rate.
- 6.24 For the liver cancer data, the conditional test yielded a positive result significant at the $0.01 < p < 0.05$ level, implying that residence near the headland is associated with an increase in the risk of liver cancer. It is important, however, to note that there may be factors other than the radioactive contamination associated with residence near the headland.
- 6.25 No such association was found for NHL using the conditional test, indicating that the excess incidence of NHL is not associated with proximity of residence to the headland.

NHS Fife report

- 6.26 At COMARE's request, a review of the medical records of each of the liver cancer patients was undertaken by NHS Fife; prior approval was obtained from the Caldicott Guardians for NHS ISD, NHS Fife and NHS Lothian. Only 8 records were available for review and, of these, one did not meet the definition of primary liver cancer. The remaining seven had pathologically proven or clinical and radiological evidence of primary liver cancer (ICD-10 Codes C22.0-22.8).
- 6.27 It should be noted that the cases from elsewhere in Scotland used to generate the expected number were not reviewed and so it is not known whether any of these may have been misclassified also, which would also affect the SIR (by increasing it). If it is assumed that the two patients for whom the records were unavailable were classified correctly, this would give the observed number of cases as 9. The corresponding SIR is 2.22, which is no longer statistically

significant at the $p=0.01$ level if the expected number of cases is assumed to be unaltered (although statistically significant at the $0.01 < p < 0.05$ level).

- 6.28 The review examined the records also for known risk factors for liver cancer, the major risk factors being infection with hepatitis B or C, evidence of heavy alcohol consumption or alcoholic liver disease and non-alcoholic steatohepatitis. Less common causes include genetic haemochromatosis, alpha-1 antitrypsin deficiency and autoimmune hepatitis (El-Serag, 2011). Of the seven cases, all had some record of having known risk factors; in six cases, this was well established. The presence of these known risk factors in the majority of the liver cancer cases is likely to have played a significant causative role.

Conclusion

- 6.29 The recent ISD analysis for the period 2000-2009 shows two types of cancer, which have been linked with exposure to radiation to some degree by previous epidemiological studies, to have statistically significantly raised incidences in the area around Dalgety Bay. The raised incidence rates have been confirmed by independent analysis.
- 6.30 It is not possible to be definitive about the cause of any particular case of cancer. Although liver cancer is classed only as '*occasionally associated with radiation*' (Boice, 2006), a statistical association with residential proximity to the headland exists and so a link with radiation cannot be excluded totally. This statistical evidence must be assessed in the light of all available information.
- 6.31 Given that it is known that the potential hazard relating to radioactive contamination at Dalgety Bay is due to the intake of radium-226, a particular pattern of cancers would be expected from epidemiological studies of those that have been exposed to radium-226, namely bone and head cancers. This signature of exposure has not been found in the cancer incidence data for Dalgety Bay, while there are indications that other (non-radiation) factors are responsible for the excess cases that have been found, and this is so for liver cancer particularly. It is thus very unlikely that the excess cases of liver cancer or of NHL during 2000-2009, or of NMSC during 1975-2002, are due to the presence of radium-226 in the area.
- 6.32 It should be noted that there is a time-lag between an individual being irradiated and the development of any consequent cancer (the latent period), which depends on the type of cancer. The above analysis, therefore, would not reflect any consequences of recent finds.

CHAPTER 7

CONCEPTUAL MODEL AND RECENT INVESTIGATIONS

Introduction

- 7.1 The history of land-use and activity on the site is reviewed in Chapter 2 of this Report, and the history of the discovery of the contamination, radiological surveys and investigations of the site is detailed in Chapter 4. This Chapter presents a review of the current state of the site and the environmental processes affecting it, with particular attention to coastal processes that may be mobilising and transporting radium sources in contaminated material from the shore onto beaches.
- 7.2 A review of the potential for radium from luminising activities to migrate in the environment was undertaken on behalf of the MoD by Baker and Toque (2005). They identify soil erosion by surface water and subsequent movement of radium-bearing particulates in surface waters as the principal means by which migration might occur, and conclude that,

'The extent to which radium will migrate in surface water from a former burial or burning site relates to a number of site-specific parameters including the accessibility of the contaminant to the surface water, the extent of vegetative cover, the intensity of rainfall and the slope of ground surface. The most likely scenario for the migration of radium in surface water would be if surface disposal of luminising ash had occurred on a poorly vegetated slope.'

- 7.3 Unfortunately, Baker and Toque (2005) did not consider erosion of coastal land by the sea as a potential scenario for mobilisation of particulate radium. That appears to be the principal mechanism at Dalgety Bay where made ground along the coastline, in which radium-bearing sources have been buried, has been eroded by waves and redistributed onto beaches and in some cases subsequently re-excavated. Exposed sources have been transported along the coastline, away from the area of the original release, by waves and tidal currents as part of the natural sediment transport process.
- 7.4 The distribution pattern of radium-bearing sources that have been recovered from beaches and foreshore at Dalgety Bay supports this interpretation. Figure 4.1 shows the distribution of items of radium-bearing materials recovered during surveys by AMEC (on behalf of the MoD) and SEPA during 2011 and 2012. No definitive map exists of finds made before this date, but it is believed that the extent of contamination found by the 2011-12 surveys is broadly similar to that of previous efforts. The 2011-12 surveys followed more thorough procedures than previously, and the much increased find rates compared with previous years may be due to this. The areas searched were mostly on the beaches, but included adjacent areas on land in some places along the coastal paths.
- 7.5 From Figure 4.1 it is immediately apparent that the distribution of finds is most dense in the area around the jetties, in the small bay marked "Mud" immediately north of the "IRB Sta", and westwards from the western promontory of this bay to the point marked 'X' (at E316360,N683320). To the west of these areas the spatial density of finds on the beach is lower, but a scatter occurs as far west as the "Outfall" in the inner, western corner of the main bay. Only a few surveys have extended beyond the "Outfall". Some isolated evidence of contamination was found in the form of count rates higher than background, but no sources have been located along the coastline beyond this point. At the other extreme of the most heavily contaminated area, the

density of sources on the beach declines immediately west of the jetty area. A scatter of finds occurred on the sandy beach that is exposed at low tide off the Headland. Two sources found in the sediments of the New Harbour were the most distant from the main contamination in this direction.

- 7.6 Many of the sources found in 2011-12 were buried within the beach sediments, some down to 0.6 m below the surface (deeper in a few cases). The search protocols introduced in 2011 were designed to locate high-activity sources buried relatively deeply as well as those on or near the surface. The find with the highest activity, 76 MBq, was buried at a depth of approximately 0.6 m, consistent with the source finds in 1990/91.
- 7.7 The pattern of finds may be conceptualised as having arisen through the erosion of the shoreline by storm waves that released material from the parts of the coast where formerly made ground is still unprotected. As will be shown below, this includes the jetty area and the “Mud” Bay to the north, whereas since 1990 the radium-bearing made ground in the Headland has been protected by a series of placements of rock materials culminating in the ‘armour’ of large blocks forming the present coastline. Thus, the densest part of the find distribution is on beaches in front of the area where radium-bearing made ground is exposed to wave attack. The peripheral part of the distribution west of point ‘X’ is most likely made up of materials which have been transported along the coast by the action of waves and tidal currents. The beach finds between point ‘X’ and the western headland of the “Mud” bay may also result from such transport, as it is not known exactly how far west the original infilling of the coast extended. The thin scatter of finds off the Headland southwest of the jetty area may similarly have resulted from wave and tidal transport, although it is likely that some material is being washed out from the interstices of the rock armour where this directly covers radium-bearing made ground. Westwards transport on this section of coast is suggested by the two finds in the New Harbour. However, the overall distribution indicates that the dominant transport direction across the Headland is likely to have been eastwards, and that the debris from the jetty area is being carried around the rocky promontory at the tip of the peninsula into the “Mud” Bay, from which transport is in turn westwards into the inner recess of the main bay.

Recent Investigations

- 7.8 In 2012, a ground penetrating radar (GPR) survey of the site was carried out by the Universities of Stirling and Belfast (Ruffell and Tyler, 2012). This identified areas of made ground, between 1 and 2 m thick, across the site. Subsequently, a similar survey was completed by a MoD sub-contractor (Zetica Ltd, 2012).
- 7.9 On the basis of the information from both GPR surveys, the MoD commissioned a contractor to undertake intrusive investigations and these were carried out in early 2013. The results were detailed in a report prepared by AMEC Environment and Infrastructure UK Ltd for the MoD (AMEC, 2013a). Figure 7.1 shows the distribution of boreholes and trial pits, together with the lines of cross sections illustrated in the AMEC report.



Figure 7.1: Map showing the locations of trial pits made in 2013, with Figure 7.7 lines of cross sections shown. (N.B. Cross sections along lines EE' and FF' are not included in this report.).

- 7.10 The work confirmed that there was a large volume of made ground containing ash and clinker behind the current rock armoured headland. On the land at the head of the jetties the made ground material was over 1 m thick. On the landward side of the coastline of the “Mud” Bay the made ground deposits were 1.5 to 2.2 m thick. On the land at point ‘X’ there was 1.5 m of made ground, thinning inland to 0.5 m. Much of the made ground material in these areas contained clinker and ash containing heterogeneously distributed radium contamination. From the presence of marine shells, there was evidence that some of the material had either been in a marine environment or had become admixed with marine sediments such as beach sand, whilst the presence of burnt sand under some deposits suggested some direct burning on the beach.
- 7.11 The made ground deposits on land either currently on or within 5 m of the coastal frontage of the peninsula have a volume of about 3,000 m³ (450 m of coastline of average thickness 1.5 m) and represent a significant potential future supply of contaminated material to the beach and marine environment. At present the deposits are not adequately protected from marine action and as a result will erode over time to repopulate the foreshore and feed further spreading of the contamination along the beaches surrounding the peninsula. Although the rock armour at the headland does offer limited protection from the most energetic incident waves, it is not fully engineered and no membrane and/or rock filter layer has been installed to prevent wash-out by waves and migration of material from within the headland onto the beach via the interstices in the rock armour. Thus, without isolation or removal, contamination will continue to be deposited on the foreshore as the coastal areas erode.
- 7.12 Figure 7.1 shows that in addition to the investigations on land, a number of trial pits were dug in the beach. The upper part of the beaches around the peninsula are mostly composed of sandy gravel containing numerous man-made objects such as fragments of brick, concrete and pottery,

as well as natural stones and sand. This coarse material forms a slope down to seaward, usually 1 or 2 m in height, with an abrupt change of grain size to sandy mud and/or silty clay at its foot. Trial pits in the upper part of the beach slope on the coast between point 'X' and the jetty area typically showed 0.5 to 1 m of made ground. Within the "Mud" bay, the made ground layer in one trial pit contained a fragment of china marked "ER 1954", indicating its deposition after that date. Towards the foot of the slope the layer reported as made ground was thinner, 0.1 to 0.2 m, and overlaid marine silt and clay. This distribution is consistent with marine erosion having cut into a layer of made ground that extended some metres to seaward of the present coastline and was formerly as thick as that on the adjacent land (i.e. 1 – 2 m as described above in paragraph 7.10), and having bevelled a new beach surface across it. The surface layer of the new beach profile would have been reworked by waves, but the deeper parts (below about 0.3 m) may represent undisturbed made ground. Some of the eroded material has been transported seawards and now forms the thinner layer resting on normal marine mud seen in the offshore trial pits.

- 7.13 West of point 'X', the grain size of the upper beach sediments changes fairly abruptly, the gravel component becoming finer over a distance of a few metres along the beach. Trial pits in the beach between point 'X' and the "Outfall" recorded only 0.2 – 0.3 m of made ground, and in some pits none was present. It seems likely that the material recorded as made ground in these pits is a reworked deposit that is rich in anthropogenic material that has been eroded from the coast line east of point 'X' and transported westwards by waves and tidal currents.
- 7.14 The distribution of made ground and radium-contaminated materials in the trial pits suggests that there may have been deposition of anthropogenic material along the whole length of coastline between point 'X' and the Headland. Subsequent marine action has eroded this material, causing recession of the coastline, with the eroded material being both re-deposited on the lower beach slope and transported along the upper beach into the inner part of the main bay (i.e. the topographic feature known as Dalgety Bay, as opposed to the town of that name). Figure 7.2 is a sketch cross section that illustrates this concept.





Figure 7.2 : Conceptual cross-section of part of the landfill and adjacent beach (not to scale).

Historical changes


- 7.15 The historical changes in the position of the coastline outlined in Chapter 2 are consistent with the concept of the current state of the site outlined in the preceding section. Historical evidence regarding the site use including aerial photographs of the coastline were reviewed by Enviro Ltd for the MoD (Enviros, 2007). These demonstrated that significant changes occurred to the coastline during the 1950s, but exact documentation of the coastline changes was not attempted.

Ordnance Survey maps normally indicate the coastline as two contour lines, for Mean High Water Spring Tide and Mean Low Water Spring Tide respectively. Neither of these corresponds exactly to the coastline as one might regard it when standing near a point where the land ends and beach begins. This latter 'morphological coastline' is usually quite obvious when one is present on the ground, but is not always easy to define as a precise line where there is a transitional zone between land and the uppermost part of a beach above high tide levels. Exact interpretation from aerial photographs can be even more problematic because what is seen on the photograph as ground uncovered by water depends on the state of the tide, weather and lighting, as well as on the height of the aircraft and quality of the photograph itself. In order to document changes over time, a series of photographic images taken many years apart must be corrected for parallax distortions and geo-referenced to a common base grid. The accuracy of this process may lead to discrepancies of several metres between different photographs in the series.

- 7.16 A study made for SEPA by the University of Stirling (Sneddon et al, 2013) has addressed these issues as far as is feasible, and presents its findings as a series of dated coastlines from 1949 to 1990. In order to minimise errors, the study treated the Headland area separately from the "Mud" bay that lies on the north-east side of the peninsula. For both areas the results were presented as a series of dated lines superimposed on a 'Google Earth' satellite image from 2009, taken at a time of high tide, and in some cases on contemporary aerial photographs also. The interpreted coastlines for 1949 and 1990 were taken as reference lines against which changes at other times could be compared. There was no significant change in the coastline of either area between 1945 and 1949, but there were substantial changes between the latter date and 1990. Since 1990 the coastline has been built outwards by emplacement of rock armour to prevent erosion of the Headland and Boatyard areas, with infilling behind a boulder screen. This has advanced the coastline by a maximum of ca.8 m in each area, as shown by a comparison of the 1990 coastline with the 2009 Google image (Figures 7.3a and 7.3b). These changes were made by the Dalgety Bay Sailing Club (see paragraph 2.19). Other parts of the coastline have not changed significantly since 1990.




a. Coastline changes in the Headland area since 1979




b. Coastline changes in the Boatyard and "Mud" bay area since 1945

Figure 7.3 : Coastline changes in two parts of the Dalgety Bay peninsula (modified from Sneddon et al, 2013).


7.17 Between 1949 and 1955, substantial changes in the coastline occurred. For the Headland area, comparison of the aerial photographs in Figures 7.4a and 7.4b shows that part of the coastline was built outwards in the triangular area to the right of the arrow in the 1949 image. In 1949 the seaward edge of this area appears to form a cliff, the line of which can still be discerned as a faint tonal change in the 1955 image. The coastline changes are directly compared in Figure 7.4c in which the two interpreted coastlines are superimposed on the 2009 Google Earth image. This extension of the coastline is clear evidence that landfilling took place between 1949 and 1955.



a. The coastline of the Headland area in 1949




b. The coastline of the Headland area in 1955




c. A comparison of the 1949 and 1955 coastlines with the Headland in 2009

Figure 7.4 : The coastline of the Headland area in 1949, 1955 and 2009 (modified from Sneddon et al, 2013).


- 7.18 The changes in the “Mud” bay area are more dramatic. The coastlines shown in Figure 7.3b indicate that this bay underwent substantial infilling between 1945 and 1990. Figures 7.5a, 7.5b and 7.5c compare the 1949, 1955, 1959 and 1990 coastlines. In 1955 a hook-shaped feature referred to by Sneddon et al. (2013) as a ‘spit’ has developed across part of the bay. This is an unfortunate choice of terminology as it is not clear whether the feature was formed by natural processes or is wholly or partly man-made. A ‘spit’ is a coastal feature that is formed naturally where a large flux of sediment is being transported along a coastline and is deposited in deeper water where the coast changes direction. The spit builds progressively outwards from the point where the original coast changed direction. The distal ends of natural spits show a characteristic curvature of beach ridges around their tip. This pattern might be represented by the curvature of lineations on the hook-like feature in Figure 7.5a, but it is hard to be sure. The tip of the feature appears to be topographically above the surrounding areas and the truncations of the lineations do not look natural. If the construction of the feature was from naturally transported sediment, it must have been deposited by currents passing through a gap between the area of rocks and sediment that lies to the east, and the tip of the land forming the main peninsula. Some of the lineations clearly pass through this gap, but the outermost ones, right at the base of the photograph, abut against the area of rocks.
- 7.19 An alternative explanation for the origin of the hook-like feature is that it was formed by deliberate dumping of material into the bay. If that were the case then the feature would have been constructed from material brought along the beach and the lineations might be interpreted as vehicle tracks. This interpretation also has problems, especially the abutment of the outermost ‘tracks’ against the area of rocks at the base of Figure 7.5a.



a. Coastline changes in the Boatyard and "Mud" bay area, 1949 to 1955




b. Coastline changes in the Boatyard and "Mud" bay area, 1955 to 1959




c. Coastline changes in the Boatyard and "Mud" bay area, 1955 to 1990

Figure 7.5 : Coastline changes in the Boatyard and "Mud" bay area (modified from Sneddon et al, 2013).


- 7.20 A third possibility, and on balance the most likely, is that the hook-like feature was deposited naturally as a natural response of the marine system to an artificially increased supply of sediment. The photos for the period between 1945 and 1949 all show a small peninsula apparently formed of sediment-covered rocks at the extreme east of the area now forming the Boatyard. The interpreted coastlines for those years show this feature as fluctuating in outline, though always of much the same general size and position. It may be that the feature represents an embryonic spit forming where the coastline turns to the north, and that the sediment needed to build and maintain it was supplied by material dumped onto the outer edge of the triangular area visible on the Headland in Figures 7.4a and 7.4b, or dumped directly onto the beach.
- 7.21 What is certain is that the hook-like feature could not have been constructed naturally without an increased supply of sediment, the likely source of which is discussed in paragraph 7.29. The northwards direction of transport implied by this is the same as that suggested by features of the present day situation on the site, namely the distribution of radioactive sources, and also the piling-up of sediment on the western sides of the jetties, with relatively lower sediment surfaces on their eastern sides.
- 7.22 Between 1955 and 1959, there were further, equally important changes in the “Mud” bay area, illustrated by Figure 7.5b. The hook shaped feature had completely disappeared by 1959 whereas the coastline of the inner part of the “Mud” bay had advanced seawards by 25 m. The area of new land in the inner bay was estimated by Sneddon et al (2013) as 3,100 m². It appears to be roughly equal to the area of the former hook-shaped feature. If these changes occurred naturally, they would have involved a major reduction or cessation of the sediment supply to the hook-shaped feature. This feature, as noted above, could have been produced by an enhanced supply of material from the Headland and jetty areas to the south. If the supply of sediment were cut off then the hook-like feature would have been unable to grow further. The same patterns of waves and tidal currents that produced the hook-like feature may then have transported the sediment westwards, depositing it against the inner coastline.
- 7.23 The changes to the “Mud” bay from 1959 to 1990 were comparatively slight. They are illustrated in Figure 7.5c. The coastline in the southern part of the bay has retreated a few metres landward whereas in the north of the bay there was a slight accretion. During the same period there were some relatively minor changes to the coastline of the Headland (see paragraph 7.30).
- 7.24 A series of cross sections based on the results of trial pit investigations made by AMEC in 2013 throw some further light on the uncertainties regarding the formation and destruction of the hook-shaped feature. Figure 7.6 shows interpolated cross sections interpreted by Nina Patton of SEPA, from the trial pit logs in AMEC (2013a), high-lighting the presence or absence of radium-contaminated ash and clinker in the made ground. The lines of these cross sections are shown in Figure 7.1. Figure 7.6a shows cross-section BB’ in the northern part of the “Mud” bay (Figure 7.1). The approximate positions of the coastline at different dates are marked and the material that accreted to this part of the coast between 1959 and 1990 was described as “ash and clinker” and was contaminated by radium. No ash was present in the pit to landward of the 1959 coastline, but this was in made ground. Unfortunately, this was the only pit in this cross section that lay in the land that was accreted between 1955 and 1959. However, it seems likely from the elevations of the deposits that the ash-free made ground in this pit was deposited before 1959.



a. Cross section through the coastline and beach along line BB' in Figure 7.1



b. Cross section through the coastline and beach along line CC' in Figure 7.1



c. Cross section through the coastline and beach along line DD' in Figure 7.1

Figure 7.6: Cross sections along the lines shown in Figure 7.1, based on the results from trial pits reported by AMEC (2013a) and reproduced from diagrams prepared by Nina Patton (SEPA) in June 2013.

- 7.25 Moving southwards, Figure 7.6b shows the cross section CC' through the central part of the bay. Between this cross section and the previous one lies the trial pit from which dated pottery from 1954 was recovered from made ground beneath the beach without radium contamination (see paragraph 7.10). If this layer of ash-free made ground was deposited at the same time as that discussed in the last paragraph, the date of deposition must have been after 1954, but before 1959.
- 7.26 Every trial pit between the 1946 coastline and the current coast in cross section CC' shows a layer of radium-contaminated ash and clinker, overlain by ash-free made ground (Figure 7.6b). The ash-and-clinker layer is exposed at the current coastline where it forms a cliff about 0.6 m high capped by a thin layer of ash-free material. It seems probable that the layers of both materials are laterally continuous. The upper, ash-free, layer of made ground overlaps the ash layer onto the land to the west. In pit TP3/3/045 the ash layer rests on fused sand. This strongly suggests that burning took place at, or very close to, this point and makes it seem unlikely that the ash layer at this location was deposited from the sea. The pit is located close to the estimated position of the 1955 coastline and so would have been on the beach at that time, but in the sea in 1946. Thus, there is evidence of a layer of ash and clinker that stretches from the 1946 coastline to at least the position of the present coastline, with further evidence of burning taking place on exposed beach sand, suggesting that some of the ash at least was produced *in situ* in the vicinity of the 1955 coastline. If the ash to seaward of this represents the transported material from the hook-shaped feature on the 1955 photograph, it implies that the feature itself must have been almost entirely composed of ashy materials, and that the ash in Section CC' was deposited not as a single layer, but as two abutting layers, one *in situ*, the other from the sea.
- 7.27 Since 1959, the coast at Section CC' (Figure 7.6b) has experienced erosion, which appears to have cut into the ash layer. This erosion may have been a source for the ashy material which has accreted to the coast in section BB' (Figure 7.6a).
- 7.28 Section DD' (Figure 7.6c) has a similar stratigraphy to Section CC' with a layer of ash and clinker with radium contamination that is overlain by ash and radium-free made ground. This section cuts through the root zone of the hook-shaped feature of 1955 (the seaward side of which is shown as "1955 spit approx."). It can be seen that this feature was likely underlain by the layer of ash and clinker. The upper layer of ash-free made ground is absent from the root zone of the hook shaped feature, but present to landward of the 1959 coastline, where it extends onto relatively high ground ca.2.5 m above the present coastline at the west end of the section. This makes it very unlikely that the ash-free made ground was deposited from the sea.
- 7.29 The evidence from the cross-sections, together with the aerial photo evidence already discussed, are consistent with the following scenario. Radium-contaminated ash and clinker was produced in the Headland area at some time before 1955 and added to the seaward edge of the triangular area visible on the 1949 photograph, as well as possibly being dumped on the beach or burned *in situ* there. Aircraft repair and reclamation commenced on the site in 1939 (paragraph 2.5), so feeding of radium-contaminated ash onto the coastline and beach may have occurred at any time during the decade prior to 1949. It is certain that some additions of material were also made between 1949 and 1955 as photographs show an accretion of land on the Headland coastline between those dates. As it was added to the beach area, the ash was entrained by coastal transport processes and carried northwards. The small peninsula or 'spit' interpreted as being present on the eastern tip of the main site from 1946 to 1950 is likely to have been composed of sands contaminated by this ash, its form being a natural response to the extra supply of artificial sediment that it added to the coastal system. Sometime between 1949/50 and 1955 the supply of sediment bearing radium-contaminated ash and clinker was increased. In the same period the

coastline of the headland was extended to seawards and the hook-like 'spit' feature developed across the "Mud" bay. On the evidence of cross-section DD' (Figure 7.6c), the hook-like feature seems mainly to have been composed of ashy material. Between 1946 and 1955 some burning to produce ashy material in situ took place on a beach that then formed the back of the bay, but now lies 20-25 m to landward of the coast (cf. cross-section CC' Figure 7.6b). In or about 1955, the supply of sediment to the hook-like feature ceased or was greatly diminished and the material composing it was washed by wave action into the bay, where it formed a layer containing radium-contaminated ash and clinker that covered the bay floor and piled up to a metre thick against the 1955 coastline (Section CC'). After this, but before 1959, a layer of ash-free made ground containing domestic artefacts dating from 1954 was deposited on the land areas adjoining the then coastline and in the bay itself. This layer was up to 2 m thick. By 1959, marine erosion had reworked some of this layer and produced beach ridges that are visible on the aerial photograph in Figure 7.5c, suggesting that the ash-free landfilling took place long enough before the 1959 photo was taken for some wave and tide action to have accomplished this.

- 7.30 Little change occurred to the Headland coastline between 1955 and 1960. After 1960 there were further changes. By 1973 small spits of land had extended into the sea in the area now grassed over in the Headland, and the coast line had accreted seawards in the jetty area. The 1973 photograph is the first to show the east jetty, the west jetty being added between 1976 and 1979. By 1979 the coast line had receded and the small spits had shifted position slightly and been reduced in size. By 1985 the first coastal armour composed of large blocks had been added to the Headland and there was little change thereafter until 1990 (Figure 7.3a). As noted above (paragraph 7.16), material and rock armour has been added to both the Headland and the Boatyard areas in the period since 1990.
- 7.31 On the coastal segment to the north of the main site, adjoining Ross Plantation, a cross-section was made by SEPA using AMEC data (Patton, 2013). This is shown in Figure 7.7 and Figure 7.1 as cross-section AA'. It is adjacent to point 'X' on Figure 4.1. It shows that the shoreface and upper beach at this point are underlain by interbedded layers of sand mixed with ash and clinker containing radium contamination. This suite of materials has probably been derived from the ashy layer in the back of the "Mud" bay, where coastal erosion has cut into it. Some of the material may derive also from further south, as investigations by SEPA and the MoD demonstrated that ashy material underlies the jetty area and is exposed to active erosion and re-deposition by waves. As already noted, in the long-term over a period of years the material released into the coastal system from this area is likely to undergo transport into the "Mud" bay and then to be carried further west into the inner part of Dalgety Bay.



7.32 So far in this Chapter only the current state of the site has been considered; the pattern of radioactive finds along beaches, the AMEC investigations of 2013 and the historical evidence of coastline changes have been used as the basis for the conceptual model of current processes and past events outlined in paragraphs 7.7, 7.14 and 7.29 and in Figure 7.2. Before drawing the evidence together it is important to consider what is known of current coastal processes on the site, and to examine the compatibility of these processes with the transport and deposition of materials that have been inferred from the evidence above.

- (a) that waves breaking in shallow water are likely to be the main mechanism for transporting the wide range of sediment grains, from coarse sand up to small boulders, that have been observed on the beaches;
- (b) that while the fetch for generating waves is greatest from the direction of the North Sea, the islands and rocks around Inchcolm that lie between due east and ESE of the site will provide shelter and refract the easterly waves round so that they approach the shore from the S to SE;
- (c) that the largest waves generated within the Firth will impact the shore from SW to ESE directions;
- (d) that shallow-water waves with significant wave heights^e of 1 to 1.25 m can be generated from time to time by the wind speeds observed in the region;

49

- (e) that such waves will move material of most sizes observed on the beach, but the larger pebbles and boulders would be moved only momentarily as waves plunged and broke over them;
 - (f) that the net direction of sediment movement along this part of the coast is from west to east; and
 - (g) that there is evidence from sedimentary structures of sediment transport from Promontory 2 north-westwards across the mouth of “Mud” bay (called “Boatyard Bay” in the report).
- 7.34 Building on these conclusions regarding processes, the Coastal Processes Report (AMEC, 2013b) states that the source of coarse debris – i.e. material up to small boulder size, including bricks or fragments of bricks and other building materials, as well as rock particles – is likely to have been erosion of the tipped debris that formed the Headland (Promontory 1) prior to the emplacement of rock armour. The report points to the disparity in beach levels across the jetties (the “groyne effect”) as further evidence for this conclusion.
- 7.35 These conclusions are in broad agreement with the evidence already considered, and do not contradict those conclusions that have already been reached from the foregoing synthesis of conceptual model and site data. It should be noted that the Coastal Processes Report omits detail on some aspects. The conclusion (c) that the largest waves may approach from the SW to SE quadrant is reasonable, but considers only the fetch available and neglects to take into account how this will combine with wind direction or the relative frequency distributions of wind speeds from different directions. The refracting effect of islands is undeniable (conclusion (b)), but in drawing conclusion (c) the Report omits the fact that the gaps between the islands are substantial and that there is 2.5 – 3 km of fetch (across shallow water) from the islands to the site. This gives an opportunity for waves generated by easterlies to pass between the islands and undergo regeneration before hitting the east-facing coasts of the site. Another omission from the Report is any mention of surges, which are an important feature of the water level regime on the east coast of Britain and could influence wave heights and the areas of the Dalgety Bay peninsula directly exposed to wave action during a surge. The impacts of these omissions are difficult to assess currently, but they may have important implications when considering the likelihood of storm wave erosion of the radium-contaminated deposits in “Mud” bay (= Boatyard Bay).
- 7.36 Further work on coastal processes, possibly including a full-scale modelling exercise, may be needed to generate valid engineering criteria for future coastal protection at the site. The nature of such protection, and of the strategy that should guide remediation, are issues yet to be decided (see paragraphs 7.45 and 7.46).

Conceptual model of the site and implications for remediation and long-term protection of the public

- 7.37 Much of the area between the modern shoreline and the 1949 coastline was filled with debris that contained radioactive sources, ash and clinker. Some of the infill was artificially deposited directly onto beaches or adjacent land areas. In some areas marine action may have been an agent in the process of infilling, following transport of materials that were dumped onto the beach within the area now designated the Headland. However, even where marine action may have been involved, there was subsequent direct deposition of landfill materials to form a surface layer. The areas on which radium-contaminated materials were deposited by one agency or another extend from the rock outcrops on the coast immediately east of the New Harbour, across the interior of the Headland, across the exposed beaches of the jetty area, around the point at the eastern extremity of the site and across the “Mud” or Boatyard bay, probably as far as the point marked ‘X’ on Figure 4.1.

- 7.38 Erosion by waves has trimmed the seaward edges of the original landfilled area, causing the high-tide waterline to recede landwards, probably by a distance of several metres in places. Material removed by waves has been deposited lower down in the area exposed at low tide, the net effect being the creation of a pebble and coarse sand beach with a profile that in some places has its upper part bevelled across radium-contaminated made ground, but in other places across made ground that lacks radium contamination. This concept is shown as a diagrammatic profile in Figure 7.2.
- 7.39 It follows from this and from the profiles in Figures 7.6 and 7.7 that radium-contaminated material is likely to be present on beaches as a veneer of reworked beach sediments less than a metre thick which in places overlies radium-contaminated material that variously represents undisturbed landfill, former beach sediments containing large amounts of ash and clinker that were dumped onto the beach, or reworked materials from the hook-shaped feature visible in "Mud" bay in 1955. Waves will disturb this veneer to varying depths depending on their direction, energy and height. Large waves coming onto a beach will tend to remove material from the upper part and deposit it further down. Storms cause erosion of the upper beach and an episodic, but progressive, erosion of the land's edge. Thus, over time the whole beach system probably increases in width at the expense of the land. Undisturbed made ground to landward of the present coastline has a volume of several thousand cubic metres and is a reservoir of contamination that is vulnerable in the long run to erosion by marine action. Such erosion would spread the contamination onto beaches and shorelines and into the marine environment, in the same manner as has already occurred.
- 7.40 The disturbance and reworking of the beach materials from time to time exposes radioactive sources at the surface, or deposits them at shallow depth where they can be detected. It is an established fact that if such sources are removed they are eventually replaced by new sources. This repopulation of the near-surface occurs when, (a) waves of all sizes and frequencies of occurrence rework the veneer of beach material and bring new sources to the surface, and (b) larger storms excavate into previously the deeper layers of undisturbed radium-contaminated material either beneath the beach veneer or at the edge of the land.
- 7.41 Parts of the coastline have been armoured with large rocks as an attempt to slow down the erosion of the land's edge. Waves may excavate finer materials from between these, and cavities have been detected in the area of landfill in front of the club-house that may be due to this process. Radioactive sources within the landfill materials behind the armour may thus be transported onto the beach in front of it.
- 7.42 Once a radioactive source has been incorporated into the beach sediments, it may be periodically buried and exposed, and then re-buried and re-exposed, unless eventually detected and removed. Some sources may break up into finer particles. On the beaches north of Ross Plantation the sediments become finer westwards from the point marked 'X' on Figure 4.1 towards the inner corner of the main bay. The more westerly part of this section of coast was not landfilled, but radioactive sources have been found in the sediments of the upper beach there. These probably result from transport along the coast of the finer fractions of the sediments that have been eroded from the landfill-backed beaches near the headland. These finer sediments have been added to the beach profiles of the inner parts of the bay.
- 7.43 All of these natural geomorphological processes are certain to continue into the future. Waves will continue to rework existing beaches, bringing radioactive sources to the surface from time to time. Larger storms will sometimes excavate down into the sub-beach landfill material, or eat laterally into the land at the top of the beaches. Even on the armoured sections of coast, erosion

by waves may continue to remove material from behind the armour. Finer sediment will be winnowed and added to beach profiles on the more sheltered coast in the inner Bay. These processes may proceed episodically and fitfully; erosion on parts of the coast may even be replaced by deposition for periods of time. But the overall trend almost certainly will be that the landfill will be relentlessly eroded in the long term. As it is already known to contain numerous radioactive sources, these will eventually be added to those already on beaches, and with further time they may become dispersed over a wider area than at present. Corrosion and attrition of sources exposed to the waves will reduce their average size in the long term, and this will lead to greater transport and dispersal, as well as creating a future supply of radioactive sources small enough to be ingested should members of the public and users of the headland come into close proximity with them.

- 7.44 The circumstances at Dalgety Bay exactly match those identified by Baker & Toque (2005) as most likely to lead to migration of radium contamination, in all respects except one. Un-vegetated surface deposits of luminising ash have been, and continue to be, exposed to erosion by surface waters and subsequent transport by moving water. The only mismatch between the circumstances envisaged by Baker and Toque (2005) and those at Dalgety Bay is that the surface water concerned is the sea, rather than rainfall or streams. Of these three agents the sea is the most powerful and persistent.

Implications for remediation strategies

- 7.45 The present distribution of contamination, and the near-certainty that the processes of erosion and re-deposition that have operated in the past will continue to spread the radioactive contamination if nothing is done, both have important implications for the design of remedial actions. Strategies for intervention are likely to fit into one of five generic categories:
- a) Do nothing apart from placing notices asking members of the public to take precautions by not entering certain areas, not digging etc.
 - b) The above plus continue to monitor beaches and land and remove sources as they are detected.
 - c) Attempt to remove the source of the contamination. Ultimately, this is the undisturbed landfill material that is present beneath land areas and beaches.
 - d) Impose barriers to migration of the sources to the beach veneer. A particular drawback to this option is that parts of the contaminated materials may be left on the seaward side of the barrier, and might have to be removed.
 - e) Create engineered structures such as breakwaters off shore with the aim of reducing incident wave energy below that required for further erosion and transport. It is highly likely, however, that such an approach would carry much greater uncertainty in its design and performance than either options (c) or (d), although in principle it could be combined with either of these.
- 7.46 Strategies (a) and (b) do not address the root of the problem, but seek merely to manage (i.e. reduce) the risk to the public by either modifying their behaviour or by keeping down the numbers of sources that might be encountered. Both strategies have a time dimension because they would have to continue for as long as the occurrence of sources on beaches persisted. It should be recognised that the overall risk of at least one member of the public experiencing harm depends on how long the situation lasts. The overall cost of strategies such as (b) that involve continued active management of the situation indefinitely will also depend on how long the

situation persists. Paragraph 7.43 makes clear that this may be assumed to be without end in the foreseeable future unless remedial action is taken.

- 7.47 Strategies (c) and (d) and, to some extent, (e) are more radical, engineered solutions to the problems presented by the current situation. Strategy (c) involves complete removal of the contaminated materials that underlie the land area as well as those that are beneath beaches. Wholesale removal of buried contamination from land areas would be disruptive and would not greatly decrease the risks to users or to the general public, if there were to be no excavation within the contaminated areas. This factor makes Strategy (d) appear attractive. However, an engineered barrier just landward of the present coastline would not enclose the contaminated 'made ground' deposits that lie beneath the beaches. These would then form a reservoir of contaminated materials that could perpetuate the problem. If a 'hard' barrier were to contain the entire extent of the contamination it would have to be built between 25 and 50 m offshore, and would have to resist marine erosion indefinitely. An alternative would be to remove any contaminated material that lay to seaward of an engineered barrier that was simpler to implement. As this would most likely involve removal of the beach down to its original natural surface prior to contamination, it would be essential to replace it with a suitably stable profile of uncontaminated materials. Failure to do so would be to invite marine erosion of the engineered barrier where it was unprotected because of the removal of the beach. Precise design of an engineered barrier and stable artificial beach profile would require further study and modelling of the site and its surroundings. Such study would be essential also if strategy (e) was adopted, on its own or in conjunction with (c) or (d). Though off-shore breakwaters might conceivably remove enough energy from the largest storm waves to prevent ongoing migration of the present coastline towards the land, this would be difficult to achieve with certainty. It is most unlikely that breakwaters could prevent reworking of beach materials and transport of exposed material along the shore, so the presently contaminated beaches would continue to act as a reservoir for the future supply of radioactive sources that might spread beyond the bounds of the present site. No matter which combination of strategies (c), (d) or (e) might ultimately be adopted, it is most important to understand that failure to take action to produce and implement an engineered solution will simply prolong the present situation and its attendant risks to the public, while also incurring on-going costs for an indefinite period into the future.

CHAPTER 8

IMPLICATIONS FOR OTHER SITES

- 8.1 It is known that there are several other sites potentially contaminated with luminising residues, some of which have been remediated.
- 8.2 Experience from previous remediation actions and the work reported above can be used to establish a set of principles for dealing with such sites.
- 8.3 If records are available, the potential amount of radium contamination and its form can be established. Of particular importance is whether luminising activities were carried out on the site, in which case it is possible that raw paint residues of potentially high radioactive content could be encountered.
- 8.4 In accordance with the recommendation of the Contaminations Working Group, surveys of the area should be carried out using a detector system with a detection limit of, or better than, 20 kBq of radium to a depth of 100 mm with 90% probability.
- 8.5 If radium contamination is detected, it should be established whether this is on/near the surface or is buried at depth. If the latter, a further consideration will be whether or not the site could be subject to erosion (as at Dalgety Bay) which could lead to unpredictable source exposure. Sites found to contain surface contamination or to be subject to erosion should be given priority.
- 8.6 Even if contamination is found to be present on a particular site, there may be no need for further action provided that it is present in a form for which there is no plausible exposure pathway for a member of the public. As an example, a concrete-covered, deeply-buried cache whose extent is well-defined and for which planning constraints to prevent excavation or building are in place, would not give rise to concern.
- 8.7 Where there are credible pathways for exposure, the associated risks and hazards should be assessed and a judgement made of the necessity for further investigation and/or remediation. Radon measurements should be made if radium contamination is suspected to lie beneath, or in close proximity to, buildings.
- 8.8 Should the circumstances make it appropriate, epidemiological studies of the locality should be instituted; particular attention should be paid to bone cancers.

CHAPTER 9

CONCLUSIONS

- 9.1 At present, there does not seem to be any comprehensive list available of sites in the UK potentially contaminated with radium. The Committee deems this to be unacceptable as it implies an unknown risk to the general population. There are instances (apparently including Dalgety Bay) where no records have been passed to purchasers of contaminated sites.
- 9.2 The radioactive contamination at Dalgety Bay appears to have arisen as a result of the luminising works carried out at Donibristle airfield and the destruction of radioactive aircraft instruments on the site.
- 9.3 Despite a letter suggesting that over 800 aircraft were dismantled on the site in 1946 alone (Wright, 1990), no records of the amount of radioactivity brought onto the site are available. It is not possible, therefore, to determine the radioactivity remaining or to estimate the longevity or magnitude of the associated hazard.
- 9.4 Monthly monitoring of the foreshore area to specific criteria (paragraph 4.26) has been in place since 2012 and is carried out by contractors on behalf of the MoD. Given that source repopulation occurs (paragraph 4.21), the monitoring and recovery programme is not preventing an encounter, but is lowering the possibility that such an encounter will occur. This is further compounded by the fact that the MOD's contractor cannot achieve the required limit of detection with current equipment on part of the beach as the local background is too high for the technique used (i.e. lower activity sources may not be detected and, hence, not removed). It is likely also that compliance with the advice given in signage etc will reduce in the coming years and this will result in an increase in the probability of an encounter. The monitoring programme covers a fixed area, but it could be expected that, as in the past, the contaminated area may spread outside that currently delineated. For these reasons, the monitoring and removal programme could be argued not to be consistent with the principle that radiation doses should be kept As Low As Reasonably Achievable (ALARA) in the medium and longer term.
- 9.5 The current monitoring programme occupies three weeks each month, requiring both capital equipment and staffing. There are associated costs for storage and disposal of the sources found since the Radioactive Substances Act (1993) (GB Parliament, 1993) restricts the disposal of such sources to unauthorised landfill sites to a maximum activity of 40 kBq per item. In addition, there are socio economic costs that the continuing presence of signage and monitoring teams is having on the local community. Thus the programme is deemed not to be cost-effective or sustainable in the longer term.
- 9.6 The undisturbed made ground to landward of the present coastline identified by the intrusive investigations has a volume of several thousand cubic metres and is a reservoir of contamination that is vulnerable in the long run to erosion by marine action.
- 9.7 The combination of soil erosion by surface water and erosion of coastal land by the sea results in a dynamic and changing environment for the release of radioactive sources. Without remediation it could be expected that this erosion will continue long-term and may result in a wider dispersion of sources.
- 9.8 Importantly, because of the heterogeneity of the radium contamination in terms of activity, solubility, size and distribution of sources within the made ground, a retrospective risk assessment based solely on previous surface finds may not provide a suitable assessment of

future risks. This was clearly demonstrated in 2011 when many more sources, some with far greater activity, were found on the beach at Dalgety Bay than had previously been known.

- 9.9 Based on concordant data from SEPA, the University of Stirling and PHE, taken in conjunction with data from the habits survey, it is concluded that the sources present at Dalgety Bay pose a potential risk to public health in terms of committed effective dose estimates.
- 9.10 Further work is necessary to determine how significant a risk exists resulting from skin contact, particularly for young children.
- 9.11 It is unlikely that there is a current significant risk posed by airborne contamination.
- 9.12 There does not appear to be a current risk from external radiation. Periodic (e.g. biennial) confirmatory monitoring should be carried out.
- 9.13 It is very unlikely that the excess cases of liver cancer or of NHL during 2000-2009, or of NMSC during 1975-2002, are due to the presence of radium-226 in the area.
- 9.14 There are three main options for remediation – removal, containment or a combination of these. In the past, for the terrestrial contamination found at Dalgety Bay, removal has been the option of choice. The current contamination, however, is more widespread and is not fixed. The Committee tends to the view that a combination of methods may be the most effective solution, but this should be subject to an evaluation process, with the required end-point being set by the regulator.

CHAPTER 10

RECOMMENDATIONS

- 10.1 Given the dynamic nature of the contamination, the indefinite continuation of a monitoring and recovery programme allied to demarcation of affected areas is not regarded as best practice and may well not be in compliance with ALARA; it would be undesirable also on economic grounds. We therefore recommend that the present monitoring and recovery programme being carried out by the MoD at Dalgety Bay should continue in the interim only until remediation can be undertaken, with the coverage and frequency to be determined by the regulator. The monitoring and recovery programme should be carried out to at least the standard specified by COMARE's Contaminations Working Group.
- 10.2 Given the long half-life of radium-226, the dynamic nature of the contamination, the number and activity of the sources being found and the unknown quantity of radium remaining, we recommend that the Scottish Government should take steps to ensure that effective remediation of the affected area is undertaken as soon as is possible.
- 10.3 We recommend also that, in conjunction with all stakeholders, an evaluation of the means of remediation should be instituted immediately, considering efficacy, practicability and cost.
- 10.4 In the interim period, since the area found to be contaminated with ash is extensive, parents should be recommended not to allow their children to dig. We recommend that this should be added to the warning signs.
- 10.5 The Committee acknowledges that the circumstances relevant to risk change over time and that periodic reviews should be undertaken. We recommend that these should include periodic re-monitoring of the external gamma dose rate of the affected areas and that a review of the epidemiological survey of cancer rates in the area should be carried out in 5-10 years time to allow for a latent period following the recent potential exposure to large numbers of sources.
- 10.6 The Committee should seek authority from Government and the devolved administrations to require information from appropriate bodies to create a UK-wide list of sites which are known to have been, or thought to be potentially, contaminated with radium. This should incorporate details of any monitoring and/or remediation known to have been carried out. The information available for each site should be evaluated and, where deemed necessary, investigation and/or remediation instituted.
- 10.7 We recommend that an accredited, UK-based measurement capability for the absolute quantification of high-activity sources should be established.
- 10.8 In the course of our investigations, it has become clear that further experimental work on skin doses should be undertaken. We recommend continued research into this issue, with particular reference to young children.

REFERENCES

- AGIR (2009). Radon and Public Health, Report of the Advisory Group on Ionising Radiation. RCE-11. HPA, Chilton.
- AMEC (2013a). Dalgety Bay Land Quality Assessment. Final Factual Investigation Report, DIO Project No.13032. Defence Infrastructure Organisation Report prepared by AMEC Environment and Infrastructure UK Limited for the Ministry of Defence, under commission DE4/4513.
- AMEC (2013b). Dalgety Bay. Coastal Processes Review. Prepared by AMEC Environment and Infrastructure Limited for the Ministry of Defence, under commission FTS3/ELMG/016.
- Baker A C and Toque C (2005). A review of the potential for radium from luminising activities to migrate in the environment. *J Rad Prot* **25**, 127-140.
- Black D (1984). Investigation of the possible increased incidence of cancer in West Cumbria. Report of the Independent Advisory Group. HMSO, London.
- Black R J, Sharp L, Finlayson A R and Harkness E F (1994). Cancer incidence in a population potentially exposed to radium-226 at Dalgety Bay, Scotland. *Br J Cancer* **69**, 140-143.
- Boice J D (2006). Ionizing radiation. In *Cancer Epidemiology and Prevention*, Third Edition. (Schottenfeld D and Fraumeni J F, eds). Oxford University Press.
- Breslin JP (2011). Dad blames radium beach for son's death. *Sunday Post* 22nd May 2011.
- Brown J and Oatway W (2012). Scoping Health Risk Assessment for Beach Users at Dalgety Bay to Support Advice to Scottish Government Given in February 2012. HPA-CRCE-040, HPA, Chilton.
- Brown R (2008). Risk Assessments for Radium Contamination at Dalgety Bay. *Radium Dalgety Bay Review*, DSTL 2008.
- Burgess P (2013). Unpublished report commissioned by SEPA.
- Camplin, B (1990). Letter to Mr G Stott, HMIPI Edinburgh, from MAFF, Fisheries Laboratory, 31 October 1990. File Ref: FLR 5445.
- Carstairs V and Morris R (1989). Deprivation, mortality and resource allocation. *Comm Med* **11**, 364-372.
- Charles M W and Gow C (2010). Skin dose from Dalgety Bay Ra-226 contamination: Dose rate measurements for ten selected samples. (Available at http://www.sepa.org.uk/radioactive_substances/publications/dalgety_bay_reports.aspx).
- Clyne F J, Garrod C J and Ly V E (2013). Radiological Habits Survey: Dalgety Bay, 2012. RL13/13. CEFAS, Lowestoft.
- COMARE (2005). Tenth Report. The incidence of childhood cancer around nuclear installations in Great Britain. HPA, Chilton.
- COMARE (2006). Eleventh Report. The distribution of childhood leukaemia and other childhood Cancers in Great Britain 1969-1993. HPA, Chilton.
- Dale, P (2012). Dalgety Bay Radium Contamination. Scottish Environment Protection Agency. (Available at www.sepa.org.uk).

- DIO (2012). Dalgety Bay Inspection, Investigation Plan. Defence Infrastructure Organisation
- El-Serag H B (2011). Hepatocellular carcinoma. *N Eng J Med* **365**, 1118-27.
- Enviros Consulting (2007). Dalgety Bay Information Review: Final Report prepared for Defence Estates under commission DE11/4416.
- Fife Council (2011). Fife Shoreline Management Plan 2011. (Available from <http://www.fifedirect.org/minisites/index.cfm?fuseaction=page.display&pageid=C040877C-B767-3F71-8454BE5167C5BC58&siteID=C03E446A-0241-A6A5-7462DD169B215841>)
- Fry S A (1998). Studies of U.S. radium dial workers; an epidemiological classic. *Radiat Res* **150**, S21-S29.
- GB Parliament (1993). Radioactive Substances Act (1993). (Available from http://www.opsi.gov.uk/acts/acts1993/ukpga_19930012_en_1)
- Gilbert R O (1987). Statistical Methods for Environmental Pollution Monitoring, Chapter 13. Van Nostrand Reinhold, New York.
- Harrison J D, Fell T P, Phipps A W, Smith T J, Ellender M, Ham G J, Hodgson A, Wilkins B T, Charles M W, Darley P J and Aydarous A Sh (2005). Health Implications of Dounreay Fuel Fragments: Estimates of Doses and Risks. HPA, Chilton. (Available at: www.sepa.org.uk/radioactive_substances/publications/dounreay_reports.aspx)
- Heaton B, Glasser F, Jones S, Bonney N, Glendinning A and Sell D (1996). Assessment of the Implications of Radium Contamination of Dalgety Bay Beach and Foreshore. Report to Scottish Office Central Research Unit. University of Aberdeen and Auris Environmental Ltd. The Scottish Office, Edinburgh.
- IAEA (2012). Worldwide proficiency test IAEA-TEL-2011-03 on the determination of natural and artificial radionuclides in soil and water.
- IARC (2012). A Review of Human Carcinogens Part D Radiation. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans, Vol 100D. IARC, Lyon.
- ICRP (1975). Report of the Task Group on Reference Man. ICRP Publication 23. *Ann ICRP* **23**, 1-480.
- ICRP (1996). Age-dependent dose to members of the public from intake of radionuclides: Part 5 compilation of ingestion and inhalation dose coefficients. ICRP Publication 72. *Ann ICRP* **26**, 1-91.
- ICRP (2002). Basic anatomical and physiological data for use in radiological protection: reference values. ICRP Publication 89. *Ann ICRP* **32**, 1-277.
- ICRP (2007). The 2007 Recommendations of the ICRP. ICRP Publication 103. *Ann ICRP* **37**, 2-4
- Jones J C and Day M J (1945). Control of hazards in the luminous dial painting industry. *Br J Ind Med* **2**, 202-208.
- Litovitz T, Whitaker N and Clark L (2010). Preventing battery ingestions: an analysis of 8648 cases, *Pediatrics* **125**, 1178-83.
- McPhail C (2011a). Minute of meeting of Dalgety Bay Particles Advisory Group, 20 December 2011. (Available from www.sepa.org.uk/radioactive_substances/dalgety_bay/particles_advisory_group.aspx)
- McPhail C (2011b). Minute of meeting of Dalgety Bay Interim Forum, 7 November 2011. (Available from www.sepa.org.uk/radioactive_substances/publications/dalgety_bay_reports.aspx)

- Milton M I L (1991). Radioactivity at Dalgety Bay, Report NRPB/VR/4/1488 for HMIPI Scotland. NRPB, Chilton.
- Nekolla E A, Kreisheimer M, Kellerer A M, Kuse-Isingschulte M, Gossner W and Speiss H (2000). Induction of malignant bone cancers in radium-224 patients: risk estimates based on improved dosimetry. *Radiat Res* **153**, 93-103.
- Nekolla E A, Walsh L and Spiess H (2010). Incidence of malignant diseases in humans injected with radium-224. *Radiat Res* **174**, 377-386.
- NRPB (1993). Estimates of late radiation risks to the UK population. *Doc NRPB* **4**(4), 15-157.
- Patton N (2013). Personal communication, SEPA.
- Pawel D J and Puskin J S (2012). US Environmental Protection Agency radiogenic risk models and projections for the US population. *Health Phys* **102**, 646-656.
- Radioactive Contaminated Land (Scotland) Regulations 2007. Scottish Statutory Instrument 2007 No 179, Stationery Office.
- RSAC (1958). Report of the Panel on Disposal of Radioactive Waste. Radioactive Substances Advisory Committee. Cabinet Office, London.
- Radioactive Substances Act 1993 Amendment (Scotland) Regulations 2011. Scottish Statutory Instrument 2011 No 207, Stationery Office.
- Ruffell A and Tyler A (2012). Ground penetrating radar survey at Dalgety Bay. Report P107 312.
- Samartzis D, Nishi N, Hayashi M, Cologne J, Cullings H M, Kodama K, Miles E F, Funamoto S, Suyama A, Soda M and Kasagi F (2011). Exposure to ionising radiation and development of bone sarcoma: new insights based on atomic-bomb survivors of Hiroshima and Nagasaki. *J Bone Joint Surg Am* **93**, 1-8.
- Scottish Government (2008). Environmental Protection Act 1990: Part IIA Contaminated Land. The Radioactive Contaminated Land (Scotland) Regulations 2007, Statutory Guidance. ISBN 978 0 7559 7058 2 (web publication only). (Available from <http://www.scotland.gov.uk/Resource/Doc/217739/0058314.pdf>)
- Smith GM (2010). What is a low dose? Letter to the Editor. *J Radiol Prot* **30**, 93-101.
- Sneddon C R, Tyler A N and Hunter P D (2013). Historic aerial photography interpretation for assisting regulatory enforcement. (Available from http://www.sepa.org.uk/radioactive_substances/publications/dalgety_bay_reports.aspx).
- Stebbins J H, Lucas H F and Steney A F (1984). Mortality from cancers of major sites in female radium workers. *Am J Ind Med* **5**, 435-459.
- Tanner R (2013). Personal communication, PHE.
- Tyler A N, Dale P, Copplestone D, Bradley S, Ewan H, McHuire C and Scott E M (2013). The radium legacy: contaminated land and the committed effective dose from the ingestion of radium contaminated materials. *Environ Int* **59**, 449-455.
- UNSCEAR (2008). Effects of Ionizing Radiation. UNSCEAR 2006 Report to the General Assembly, United Nations, New York.
- Watson S J, Jones A L, Oatway W B and Hughes JS (2005). Ionising Radiation Exposure of the UK Population: 2005 Review. HPA-RPD-001. HPA, Chilton.

Wilkins B (2008). Email to SEPA from HPA.

Wright I W W (1990). Letter to Lord James Douglas-Hamilton from HMIPI, Edinburgh. File DD9/724, National Archives of Scotland, Edinburgh.

Zetica Ltd (2012). Ground check report commissioned by AMEC. Appendix C of DIO (2012).

ACKNOWLEDGEMENTS

The Committee is grateful to the Scottish Environment Protection Agency for permission to use data from documents published on their website and to Public Health England for the dosimetry calculations referred to in paragraph 5.36.

APPENDIX A

GLOSSARY & ABBREVIATIONS

ACCRETION

Increase in size – particularly of an area of land as a result of deposition of material from the sea.

ALIENATED SITE

An area of land which was previously owned by, occupied by or the responsibility of the Ministry of Defence.

BACKGROUND RADIATION

Radiation that comes from naturally occurring radioactive material in the ground and from cosmic rays irradiating the earth from outer space. The UK average dose from natural background radiation is 2.2 mSv per year: regional averages range from 1.5 - 7.5 mSv per year.

BECQUEREL (Bq)

A unit of radiation equal to one disintegration per second. Discharges are normally expressed in:

Megabecquerels (MBq) – one million Bq

Gigabecquerels (GBq) – one thousand million Bq

Terabecquerels (TBq) – one million million Bq

COMBUSTIBLE

Capable of being burnt.

EPIDEMIOLOGY

The study of death or illness in a population and their relation to factors regarded as possible causes.

EROSION

The removal of solid material on the coastline by the action of the sea.

GRAY (Gy)

The international (SI) unit of absorbed dose. 1 Gray is equivalent to 1 joule of energy absorbed per kg of matter such as body tissue.

HALF-LIFE

The time a radioactive a given quantity of radioactive substance takes to lose half its radioactivity through decay. It is a consequence of the independence of atomic disintegrations that this quantity does not depend on the initial quantity, leading to exponential decay.

HAZARD

A property that in particular circumstances could lead to harm, for example, exposure to radiation leading to damage to an individual's health.

INCIDENCE

The number of instances of illness commencing, or of persons falling ill, during a given period in a specified population. More generally, the number of new events, e.g. new cases of disease in a defined population, within a specified period of time. The term incidence is sometimes used to denote

'incidence rate', i.e. the number of cases divided by the (average) number at risk in the relevant time period.

INFILL

The filling of existing spaces, for example with material deposited by the sea or placed to reinforce a barrier.

INGESTION

Intake of food or drink into the body.

INHALATION

Breathing in of vapours or gases.

IONISING RADIATION

Radiation that is sufficiently energetic to remove electrons from atoms in its path. In human or animal exposures ionising radiation can result in the formation of highly reactive particles in the body which can cause damage to individual components of living cells and tissues.

ISOTOPE

One of a number of alternative forms of given chemical element with the same atomic number but differing numbers of neutrons

LUMINISING

Making something luminous, particularly painting instruments with light-emitting paint.

NON-HODGKIN LYMPHOMA (NHL)

A group of lymphomas that differ in important ways from Hodgkin lymphoma and are classified according to the microscopic appearance of the cancer cells. In children, NHL and leukaemias are often combined due to historical difficulties in making diagnostic distinctions.

RADIATION

The emission and propagation of energy by means of rays or waves or sub-atomic particles.

RADIOISOTOPE

A radionuclide that is also one of a number of isotopes of a given element

RADIONUCLIDE

A type of atomic nucleus which is unstable and which may undergo spontaneous decay to another atom by emission of ionising radiation (usually alpha, beta or gamma).

RAMSAR SITES

International important wetland sites, including some tidal waters, designated under the Ramsar Convention signed in Ramsar, Iran in 1971.

RELATIVE RISK (RR)

The ratio of the risk of disease or death among those exposed to a potential hazard to the risk among those not exposed to the hazard.

RISK

A combination of the probability, or frequency, of occurrence of a defined hazard and the magnitude of the consequences of the occurrence. (See *HAZARD* and *RELATIVE RISK*.)

Risk is sometimes taken to mean the probability that an event will occur, e.g. that an individual will become ill or die within a stated period of time or age. Risk is also used as a non-technical term encompassing a variety of measures of the probability of a (generally) unfavourable outcome.

ROCK ARMOUR

Rocks and other material used to protect a shoreline from erosion, usually of an irregular form that dissipates the energy of the sea by permitting some permeability.

SIEVERT (Sv)

The international (SI) unit of effective dose obtained by weighting the equivalent dose in each tissue in the body with ICRP recommended tissue weighting factors and summing over all tissues. Because the sievert is a large unit, effective dose is commonly expressed in millisieverts (mSv) – i.e. one-thousandth of one sievert. The average annual radiation dose received by members of the public in the UK is ~2.7 mSv.

APPENDIX B

REPORTS OF THE COMMITTEE ON MEDICAL ASPECTS OF RADIATION IN THE ENVIRONMENT

<i>COMARE Fourteenth Report.</i>	Further consideration of the incidence of childhood leukaemia around nuclear power plants in Great Britain. HPA, Chilton, May 2011.
<i>COMARE Thirteenth Report.</i>	The health effects and risks arising from exposure to ultraviolet radiation from artificial tanning devices. HPA, Chilton, June 2009.
<i>COMARE Twelfth Report.</i>	The impact of personally initiated X-ray computed tomography scanning for the health assessment of asymptomatic individuals. HPA, Chilton, December 2007.
<i>COMARE Eleventh Report.</i>	The distribution of childhood leukaemia and other childhood cancer in Great Britain 1969-1993. HPA, Chilton, July 2006.
<i>COMARE Tenth Report.</i>	The incidence of childhood cancer around nuclear installations in Great Britain. HPA, Chilton, June 2005.
<i>COMARE Ninth Report.</i>	Advice to Government on the review of radiation risks from radioactive internal emitters carried out and published by the Committee Examining Radiation Risks of Internal Emitters (CERRIE). NRPB, Chilton, October 2004.
<i>COMARE Eighth Report.</i>	A review of pregnancy outcomes following preconceptional exposure to radiation. NRPB, Chilton, February 2004.
<i>COMARE Seventh Report.</i>	Parents occupationally exposed to radiation prior to the conception of their children. A review of the evidence concerning the incidence of cancer in their children. NRPB, Chilton, August 2002.
<i>COMARE & RWMAC* Joint Report.</i>	Radioactive contamination at a property in Seascale, Cumbria. NRPB, Chilton, June 1999.
<i>COMARE Sixth Report.</i>	A reconsideration of the possible health implications of the radioactive particles found in the general environment around the Dounreay nuclear establishment in the light of the work undertaken since 1995 to locate their source. NRPB, Chilton, March 1999.
<i>COMARE Fifth Report.</i>	The incidence of cancer and leukaemia in the area around the former Greenham Common Airbase. An investigation of a possible association with measured environmental radiation levels. NRPB, Chilton, March 1998.
<i>COMARE Fourth Report.</i>	The incidence of cancer and leukaemia in young people in the vicinity of the Sellafield site, West Cumbria: further studies and an update of the situation since the publication of the report of the Black Advisory Group in 1984. Department of Health, London, March 1996.

* Radioactive Waste Management Advisory Committee.

- COMARE & RWMAC* Joint Report.* Potential health effects and possible sources of radioactive particles found in the vicinity of the Dounreay nuclear establishment. HMSO, London, May 1995.
- COMARE Third Report.* Report on the incidence of childhood cancer in the West Berkshire and North Hampshire area, in which are situated the Atomic Weapons Research Establishment, Aldermaston and the Royal Ordnance Factory, Burghfield. HMSO, London, June 1989.
- COMARE Second Report.* Investigation of the possible increased incidence of leukaemia in young people near the Dounreay nuclear establishment, Caithness, Scotland. HMSO, London, June 1988.
- COMARE First Report.* The implications of the new data on the releases from Sellafield in the 1950s for the conclusions of the Report on the Investigation of the Possible Increased Incidence of Cancer in West Cumbria. HMSO, London, July 1986.

* Radioactive Waste Management Advisory Committee.

APPENDIX C

COMMITTEE ON MEDICAL ASPECTS OF RADIATION IN THE ENVIRONMENT

CHAIRMAN

Professor A Elliott BA PhD DSc CPhys FInstP FIPEM
University of Glasgow

MEMBERS

Dr J Bithell BA MA DPhil
Childhood Cancer Research Group, Oxford

Dr P Darragh MD PhD MSc FRCP FFPHM
Public Health Agency for Northern Ireland, Belfast

Professor W Evans MA PhD FInstP FIPEM HonMRCR
University Hospital of Wales, Cardiff

Professor S Hodgson BM BCh DM FRCP
Department of Clinical Development Sciences, St George's University of London

Professor P Hoskin
Mount Vernon Cancer Centre, Northwood

Dr B Howard MBE
Centre for Ecology and Hydrology, Lancaster Environment Centre

Professor M Kadhim PhD
Department of Biological and Medical Sciences, Oxford Brookes University, Oxford

Professor P Marsden MSc PhD FSRP MIPEM MInstP CRadP
UCL Hospitals NHS Foundation Trust, London

Dr G Maskell MA FRCP FRCR
Department of Radiology, Royal Cornwall Hospital, Truro

Professor S McKeown MA PhD FSB CBiol
School of Biomedical Sciences, University of Ulster, Coleraine

Dr T Nunan MD FRCP FRCR
Nuclear Medicine Physician

Dr M Pearce BSc MSc PhD
Institute of Health and Society, Newcastle University, Newcastle upon Tyne

Mr I Robinson BSc CRadP FSRP FNucl

Consultant on nuclear and radiation safety (formerly HM Superintending Inspector of Nuclear Installation, Office for Nuclear Regulation)

Professor R Taylor MA, FRCPE, FRCP, FRCR
School of Medicine, Swansea University

Professor R Wakeford BSc PhD CSci CPhys FInstP CStat CEng MINuE CRadP FSRP
Dalton Nuclear Institute, University of Manchester

Professor P Warwick BA MSc PhD DSc CChem FRSC
Centre for Environmental Studies, Loughborough University

Professor Catharine West BA PhD
University of Manchester

SECRETARIAT

Mr S Ebdon-Jackson BSc MSc FRCR HonFRCP (Scientific)

Dr E Petty BSc PhD (Scientific)

Dr K Broom BSc DPhil CBiol FSB (Scientific)

Ms K Stonell (Minutes)

Ms J Humphries (Administrative)

ASSESSORS IN ATTENDANCE REPRESENTING THE FOLLOWING ORGANISATIONS

Department for Children, Schools and Families
Department for Communities and Local Government
Department of Energy and Climate Change
Department of the Environment, Food and Rural Affairs
Department of Health
Department of Health, Social Services and Public Safety (Northern Ireland)
Department for Innovation, Universities and Skills
Environment Agency
Food Standards Agency
Health and Safety Executive
Medical Research Council
Ministry of Defence
Nuclear Decommissioning Authority
Office for National Statistics
Public Health England
Public Health and Intelligence, NHS National Services Scotland
Scottish Environment Protection Agency
Scottish Government
Welsh Government

WRITING GROUP MEMBERSHIP

Professor A Elliott BA PhD DSc CPhys FInstP FIPEM
University of Glasgow

Professor T C Atkinson BSc PhD
Department of Earth Sciences, University College London

Dr J Bithell BA MA DPhil
Childhood Cancer Research Group, Oxford

Mr R Black MA
National Information and Intelligence, NHS National Services Scotland

Dr T Bruce BSc MSc PhD
School of Engineering, University of Edinburgh

Professor R Wakeford BSc PhD CSci CPhys FInstP CStat CEng MINucE CRadP FSRP
Dalton Nuclear Institute, University of Manchester

APPENDIX D
DECLARATION OF MEMBERS' INTERESTS
CODE OF PRACTICE

1. Introduction

This code of practice guides members of COMARE as to the circumstances in which they should declare an interest in the course of the Committee's work. To avoid any public concern that commercial interests of members might affect their advice to Government, Ministers have decided that information on significant and relevant interests of members of its advisory committees should be on the public record. The advice of the Committee frequently relates to matters which are connected with the radiation industry generally and, less frequently, to commercial interests involving radioactivity. It is therefore essential that members should comply with the code of practice which is set out below.

2. Scope and definitions

This code applies to members of COMARE and its subcommittees, subgroups, working groups and working parties which may be formed.

For the purposes of this code of practice, the 'radiation industry' means:

- (a) companies, partnerships or individuals who are involved with the manufacture, sale or supply of products processes or services which are the subject of the Committee's business. This will include nuclear power generation, the nuclear fuel reprocessing industry and associated isotope producing industries, both military and civil and also medical service industries;
- (b) trade associations representing companies involved with such products;
- (c) companies, partnerships or individuals who are directly concerned with research or development in related areas;
- (d) interest groups or environmental organisations with a known interest in radiation matters.

This excludes government departments, professional bodies, international organisations and agencies.

It is recognised that an interest in a particular company or group may, because of the course of the Committee's work, become relevant when the member had no prior expectation this would be the case. In such cases, the member should declare that interest to the Chairman of the meeting and thereafter to the Secretariat.

In this code, 'the Department' means the Department of Health, and 'the Secretariat' means the secretariat of COMARE.

3. Different types of interest – definitions

The following is intended as a guide to the kinds of interests which should be declared. Where a member is uncertain as to whether an interest should be declared they should seek guidance from the Secretariat or, where it may concern a particular subject which is to be considered at a meeting, from the Chairman at that meeting. Members of the Committee and the Secretariat are under no obligation to search out links between one company and another, for example where a

company with which a member is connected has a relevant interest of which the member is not aware and could not reasonably be expected to be aware.

If members have interests not specified in these notes but which they believe could be regarded as influencing their advice they should declare them to the Secretariat in writing and to the Chairman at the time the issue arises at a meeting.

3.1 Personal interests

A personal interest involves current payment to the member personally. The main examples are:

- (a) *Consultancies and/or direct employment*: any consultancy, directorship, position in or work for the radiation industries which attracts regular or occasional payments in cash or kind.
- (b) *Fee-paid work*: any work commissioned by those industries for which the member is paid in cash or kind.
- (c) *Shareholdings*: any shareholding in or other beneficial interest in shares of those industries. This does not include shareholdings through unit trusts or similar arrangements where the member has no influence on financial management.
- (d) *Membership or affiliation*: any membership role or affiliation that the member or close family member has to clubs or organisations with an interest or involvement in the work of the Department. This will not include professional bodies, organisations and societies.

3.2 Non-personal interests

A non-personal interest involves current payment which benefits a department to which a member is responsible, but is not received by the member personally. The main examples are:

- (a) *Fellowships*: the holding of a fellowship endowed by the radiation industry.
- (b) *Support by industry*: any payment, other support or sponsorship by the radiation industry which does not convey any pecuniary or material benefit to a member personally, but which does benefit their position or department, e.g.:
 - (i) a grant from a company for the running of a unit or department for which a member is responsible;
 - (ii) a grant or fellowship or other payment to sponsor a post or a member of staff in a unit or department for which a member is responsible. This does not include financial assistance for students, but does include work carried out by postgraduate students and non-scientific staff, including administrative and general support staff;
 - (iii) the commissioning of research or work by, or advice from, staff who work in a unit or department for which a member is responsible.
- (c) *Support by charities and charitable consortia*: any payment, other support or sponsorship from these sources towards which the radiation industry has made a specific and readily identifiable contribution. This does not include unqualified support from the radiation industry towards the generality of the charitable resource.
- (d) *Trusteeships*: where a member is trustee of a fund with investments in the radiation industry, the member may wish to consult the Secretariat about the form of declaration which would be appropriate.

3.3 Specific interests

A specific interest relates explicitly to the material, product, substance or application under consideration by the Committee.

A member must declare a personal, specific interest if they currently receive a payment, in any form, for any significant fundamental development work undertaken previously or at this time, on a material, product or substance or its application under consideration. This will include the production of radioactive substances and devices designed to use ionising or non-ionising radiation for diagnostic, treatment or other purposes.

A member must declare a non-personal, specific interest if they are aware that the department to which they are responsible currently receives payment for significant fundamental development work undertaken previously or at this time, on a material, product or substance or its application under consideration but they have not personally received payment for that work in any form. This will include the production of radioactive substances and devices designed to use ionising or non-ionising radiation for diagnostic, treatment or other purposes.

3.4 Non-specific interests

A non-specific interest relates to a company or associated material, product, substance or application, but not to the specific material, product, substance or application under consideration by the Committee.

A member must declare a personal, non-specific interest if they have a current personal interest with a material, product, substance or application from a particular company, which does not relate specifically to the material, product, substance or application from that company under consideration.

A member must declare a non-personal, non-specific interest if they are aware that the department to which they are responsible is currently receiving payment from the company concerned which does not relate specifically to a material, product, substance or application under discussion.

If a member is aware that a material, product, substance or their application under consideration is or may become a competitor of a material, product or substance manufactured, sold or supplied by a company in which the member has a current personal interest, they should declare their interest in the company marketing the rival material, product or substance.

Members are under no obligation to seek out knowledge of such work done for or on behalf of the radiation industry within departments to which they are responsible if they would not reasonably expect to be informed. This applies to all non-personal, specific and non-specific interests.

4. Declaration of interests

4.1 Declaration of interests to the Secretariat

Members should inform the Secretariat in writing when they are appointed of their current personal and non-personal interests and annually in response to a Secretariat request. Only the name of the company (or other body) and the nature of the interest is required; the amount of any salary, fees, shareholding, grant, etc, need not be disclosed. An interest is *current* if the member has a continuing financial involvement with the industry, e.g. if they hold shares in a radiation company, have a consultancy contract, or if the member or the department to which they are responsible is in the process of carrying out work for the radiation industry. Members are asked to inform the Secretariat at the time of any change in their personal interests, and may be invited to complete a form of declaration when required. It would be sufficient if changes in non-

personal interests are reported at the next annual declaration following the change. (Non-personal interests involving less than £5000 from a particular company in the previous year need not be declared.)

The register of interests should be kept up-to-date and be open to the public.

4.2 Declaration of interests at meetings and participation by members

Members are required to declare relevant interests at Committee meetings and to state whether they are personal or non-personal interests. The declaration should include an indication of the nature of the interest.

(a) If a member has a current (personal or non-personal) interest in the business under discussion, they will not automatically be debarred from contributing to the discussion subject to the Chairman's discretion. The Chairman will consider the nature of the business under discussion and of the interest declared (including whether it is personal or non-personal) in deciding whether it would be appropriate for the relevant member to participate in the item.

(b) If a member has an interest which is not current in the business under discussion, this need not be declared unless not to do so might be seen as concealing a relevant interest. The intention should always be that the Chairman and other members of the Committee are fully aware of relevant circumstances.

A member, who is in any doubt as to whether they have an interest which should be declared, or whether to take part in the proceedings, should ask the Chairman for guidance. The Chairman has the power to determine whether or not a member with an interest shall take part in the proceedings.

If a member is aware that a matter under consideration is or may become a competitor of a product, process or service in which the member has a current personal interest, they should declare the interest in the company marketing the rival product. The member should seek the Chairman's guidance on whether to take part in the proceedings.

If the Chairman should declare a current interest of any kind, they should stand down from the chair for that item and the meeting should be conducted by the Deputy Chairman or other nominee if the Deputy Chairman is not there.

Members' declarations of interests – 2013

Member	Company	Personal Interest	Company	Non-Personal Interest
Dr J Bithell		None		None
Dr P Darragh		None		None
Prof A Elliott		None		None
Prof W Evans		None		None
Prof S Hodgson		None		None
Prof P Hoskin		None		None
Dr B Howard		None		None
Prof M Kadhim				
Prof P Marsden		None		None
Dr G Maskell		None		None
Prof S McKeown				
Dr T Nunan		None		None
Dr M Pearce		None		None
Prof R Taylor		None		None
Mr I Robinson	1. AMEC	Consultancy		None
Prof R Wakeford	1. Sellafield Ltd 2. Compensation Scheme for Radiation-linked Diseases 3. Canadian Nuclear Safety Commission 4. Augean	Consultancy Consultancy Contract Contract		
Prof P Warwick	1. Enviro Ltd 2. Sellafield Ltd/Golder 3. NNL/LLWR Ltd	Director & Shareholder Contract Consultancy	NDA	Grants
Prof C West		None		None