

Ecological and socio-economic factors affecting extinction risk in parrots

George Olah¹ · Stuart H. M. Butchart^{2,3} · Andy Symes² ·
Iliana Medina Guzmán⁴ · Ross Cunningham¹ ·
Donald J. Brightsmith⁵ · Robert Heinsohn¹

Received: 10 August 2015 / Revised: 25 November 2015 / Accepted: 6 December 2015 /
Published online: 21 January 2016
© Springer Science+Business Media Dordrecht 2016

Abstract Parrots (*Psittaciformes*) are among the most threatened bird orders with 28 % (111 of 398) of extant species classified as threatened under IUCN criteria. We confirmed that parrots have a lower Red List Index (higher aggregate extinction risk) than other comparable bird groups, and modeled the factors associated with extinction risk. Our analyses included intrinsic biological, life history and ecological attributes, external anthropogenic threats, and socio-economic variables associated with the countries where the parrot species occur, while we controlled for phylogenetic dependence among species. We found that the likelihood of parrot species being classified as threatened was less for species with larger historical distribution size, but was greater for species with high forest dependency, large body size, long generation time, and greater proportion of the human population living in urban areas in the countries encompassing the parrots' home ranges. The severity of extinction risk (from vulnerable to critically endangered) was positively related to the per capita gross domestic product (GDP) of the countries of occurrence, endemism to a single country, and lower for species used as pets. A disproportionate

Communicated by Stuart Pimm.

Electronic supplementary material The online version of this article (doi:[10.1007/s10531-015-1036-z](https://doi.org/10.1007/s10531-015-1036-z)) contains supplementary material, which is available to authorized users.

✉ George Olah
george.olah@anu.edu.au

¹ Fenner School of Environment and Society, The Australian National University, Frank Fenner Building No 141, Canberra, ACT 0200, Australia

² BirdLife International, David Attenborough Building, Pembroke St, Cambridge CB23QZ, UK

³ Department of Zoology, University of Cambridge, Downing St, Cambridge CB23EJ, UK

⁴ Department of Ecology, Evolution and Genetics, Research School of Biology, The Australian National University, Canberra, Australia

⁵ Department of Veterinary Pathobiology, Schubot Exotic Bird Health Center, Texas A&M University, College Station, TX, USA

number of 16 extinct parrot species were endemic to islands and single countries, and were large bodied, habitat specialists. Agriculture, hunting, trapping, and logging are the most frequent threats to parrots worldwide, with variation in importance among regions. We use multiple methods to rank countries with disproportionately high numbers of threatened parrot species. Our results promote understanding of global and regional factors associated with endangerment in this highly threatened taxonomic group, and will enhance the prioritization of conservation actions.

Keywords Extinction risk · Anthropogenic threat · *Psittaciformes* · IUCN · Comparative analysis · Socioeconomics

Introduction

The current worldwide decline in biodiversity requires urgent action (Butchart et al. 2010; Pimm et al. 2014; Tittensor et al. 2014), but determining appropriate actions is often impeded by poor understanding of the diverse causes of decline and extinction across different taxa and regions of the world (Spray and McGlothlin 2003). The factors that determine a species' extinction risk fall into two main categories: (1) intrinsic biological attributes of species, and (2) exposure to external anthropogenic threats (Fisher et al. 2003). Although exposure to threats is the ultimate cause of extinction, a species' biology often determines how sensitive it is to different threat types. Particular biological attributes can allow populations to recover rapidly from depletion and can offer a degree of resilience from external threats (Cardillo et al. 2004). Biological attributes that have been shown to decrease the risk of extinction include large geographical distribution, high population density, lower trophic level, and various life history traits including high reproductive rate, small body size, short generation length, and a low degree of habitat specialization (Owens and Bennett 2000; Purvis et al. 2000).

The order *Psittaciformes* (parrots) contains 398 extant species divided into 3 families (*Psittacidae* 374, *Cacatuidae* 21, *Strigopidae* 3 species) of which 111 (28 %) are classified by BirdLife International (2014) as threatened on the IUCN Red List [i.e. in the categories of Critically Endangered, Endangered or Vulnerable (IUCN 2014)]. Sixty species are classified as Near Threatened and the rest (227 species) are considered as Least Concern. *Psittaciformes* is the fourth largest bird order after *Passeriformes* (5913 spp., 10 % threatened), *Caprimulgiformes* (593 spp., 9 % threatened), and *Piciformes* (484 spp., 7 % threatened), but it contains the second highest number of threatened bird species after *Passeriformes* (containing 611 threatened species). Data from BirdLife International underpinning assessments for the IUCN Red List show that 56 % of all parrot species are in decline, 35 % are stable and only nine per cent have increasing populations (IUCN 2014).

Parrots are currently native to 124 countries between the latitudes N 35° and S 56°, but are mainly distributed in tropical and subtropical habitats of the southern hemisphere (Fig S1). They have been popular as pets throughout human history, probably due to their colourful appearance, reputedly high intelligence, and remarkable ability to mimic various sounds including the human voice (Grahl 1990). This popularity has led many species to be captured from the wild for pets. Although captive breeding has proven possible for many species (Arndt 2007; Tella and Hiraldo 2014), *Psittaciformes* remains the most common avian order reported in the wildlife trade (Bush et al. 2014).

Parrots exhibit many of the traits known to be associated with extinction: for example, many are large-bodied and slow-breeding (e.g. macaws and cockatoos), and 70 % are ecologically specialized (e.g. forest species) (Forshaw 2011). The main anthropogenic threats to parrots are habitat loss, degradation and fragmentation driven by agriculture, logging and commercial and residential development, and hunting and trapping (Beissinger et al. 1992; BirdLife International 2014; Laurance et al. 2002; Snyder et al. 2000). The socio-economic status of a country can determine the severity of anthropogenic threats affecting the species living there. For example, human population density was found to be closely related to the proportion of threatened bird species per country (Kerr and Currie 1995). Human activities such as poaching, hunting, and land clearing can be poverty driven in certain developing areas (Blaikie and Jeanrenaud 1997), but the effects of socio-economic drivers on the extinction risk of parrots have never been examined.

Although parrots have long been considered a group of especially high conservation concern (Pasquier 1980), in-depth quantitative analysis of the nature and trends in threats across the order has been lacking. Reviews to date have assessed particular threats (Beissinger and Bucher 1992; Harris et al. 2012; Laurance et al. 2009; Müller 2000; Newton 1994; Pain et al. 2006; Pires 2012), focused on individual species (Baker-Gabb 2011; Holdsworth and Starks 2006; Snyder et al. 2000; Webster et al. 2003) or assessed general trends in parrot conservation without comparative analysis (Collar 1997, 2000; Collar and Juniper 1992; Martin et al. 2014). Jones et al. (2006) advocated decision tree analysis for assessing extinction risk and used the IUCN Red List data of parrots from 2000 as an example group, but did not cover extrinsic extinction threats or socio-economic traits, and nor did they control for phylogenetic dependence of data. Our aim is to use comparative analyses to enhance understanding of the extrinsic, intrinsic, global and regional factors associated with endangerment and extinction in this highly threatened taxonomic group.

Methods

Red List Index

We examined trends in extinction risk during 1988–2012 for parrots (*Psittaciformes*) and other high profile ecological groups with similar numbers of species using Red List Indices (Butchart et al. 2007). These groups included waterbirds, seabirds, and raptors, each of which comprises multiple orders (see online supplementary methods). Other groups included pigeons (*Columbiformes*) and gamebirds (*Galliformes*). IUCN Red List criteria (e.g. absolute population size, range size, rates of decline, etc.) are used to assign species to Red List categories of relative extinction risk, and the Red List Indices are calculated from changes between these categories. Red List Indices for sets of species are based on the number of species in each category, and the number moving between categories owing to genuine improvement or deterioration in status, i.e. increases or decreases in population size, population trends, extent of occurrence, etc. that are of sufficient magnitude to cross the thresholds for lower or higher Red List categories (Butchart et al. 2004, 2005, 2007).

Database compilation

We assembled a database of the biological and geographic attributes of all 398 extant parrot species using the 2014 version of BirdLife International and IUCN's database which underpins the IUCN Red List assessments for birds on the BirdLife Data Zone (BirdLife

Table 1 Explanatory variables used in the statistical models

Group	Variable	Source of data	Description	N	Median
A	Median latitude	BirdLife International and NatureServe (2014)	Absolute value of the median latitude of the species distribution (decimal degree)	398	9.5
	Historical distribution size*	BirdLife International and NatureServe (2014)	Area of the historical distribution size of the species (km ²)	398	177,782
	Mean altitude	BirdLife International (2014)	Mean elevation of the species distribution range (m)	322	322
	Region	Olson et al. (2001)	Distribution of the species: (1) Neotropics, (2) Afrotropics, (3) Indomalaya, (4) Australasia/Oceania	398	
	Island endemism	BirdLife International (2014)	If endemic to islands smaller than 110,000 km ² (yes/no)—Large islands were not considered as islands (New Guinea, Borneo, Madagascar, Sumatra, Celebes/Sulawesi, New Zealand, Java)	398	
B	Body size*	Forshaw (2011)	Length of the species (cm)	398	26
	Number of habitats	BirdLife International (2014)	Total number of habitats utilised by species (importance: major, marginal, or suitable)	398	3
	Migrant status	BirdLife International (2014)	(0) Not a Migrant, (1) Nomadic, (2) Altitudinal Migrant, (3) Full Migrant	398	
	Main diet	Juniper and Parr (2003)	(1) Frugivore (fruits, vegetable matters, leaves, fungi, lichens), (2) Granivore (grass seeds), (3) Nectarivore (nectar, pollen, flowers), (4) Tree seeds (hard seeds, acorns, nuts, cone seeds), (5) Specialist	398	
	Social flocking	Forshaw (2011), Arndt (2007)	Flock size in the non-breeding season: (0) No flocks (alone or pairs), (1) Small flocks (up to 20), (2) Large flocks (more than 20)	397	
	Colony nesting	Forshaw (2011)	Yes/no	386	
	Nesting tree type	Forshaw (2011), Juniper and Parr (2003)	(1) hardwood trees and their branches; (2) palm trees; (3) other nest types (e.g. termite mounds, epiphytes, moss, burrows, grass, etc.)	322	
	Forest dependency	BirdLife International (2014)	(0) Non-forest, (1) Low, (2) Medium, (3) High. Scored from published and unpublished information on the ecology of each species	398	
	Generation time	BirdLife International (2014)	Mean generation length in years	398	6

Table 1 continued

Group	Variable	Source of data	Description	N	Median
C	Captive breeding	Arndt (2007)	Yes/no	397	
	Used for pets	BirdLife International (2014)	Yes/no	398	
	Used for food	BirdLife International (2014)	Yes/no	398	
	Used for accessories	BirdLife International (2014)	Yes/no	398	
	Used for sport	BirdLife International (2014)	Yes/no	398	
D	Single country endemic	BirdLife International (2014)	If endemic to one country (yes/no)	398	
	Per capita GDP	CIA (2013), IMF (2013)	Gross domestic product based on purchasing-power-parity (PPP) per capita in USD	397	10,148
	Industrial production growth rate	CIA (2013)	Mean percentage of the industrial production growth rate of countries where species is extant (%)	378	4.1
	Unemployment rate	CIA (2013), IMF (2013)	Mean percentage of the labour force that is without jobs of countries where species is extant (%)	395	6.6
	Human population density	FAOSTAT (2013), CIA (2013), IMF (2013)	Mean population density of countries where species is extant (people/1000Ha)	398	315
	Urban population	FAOSTAT (2013)	Mean percentage of population living in urban areas of countries where species is extant (%)	398	68.3
	Human population growth rate	CIA (2013)	Mean population growth rate of countries where species is extant (%)	398	1.1
	Agriculture area	FAOSTAT (2013)	Mean percentage of agricultural area of countries where species is extant (%)	398	31.3

Variables marked with an asterisk were log_e transformed

International 2014) and IUCN Red List website (IUCN 2014). Table 1 shows all explanatory variables (including sources) used in our statistical models of threat status. The online supplementary methods provide further details of the derivation of values in the database.

We used historical distribution size instead of current distribution sizes to test if species with originally small distributions are more prone to be endangered, and in order to avoid possible circularity as current extent of occurrence is a parameter used in the IUCN Red

List criteria. Historical distribution is mapped using the same sources as for contemporary distribution, adding areas with historical records prior to 1980 for which the species is now judged to be extirpated owing to lack of recent records despite searches, lack of suitable habitat and informed by expert judgment (BirdLife International and NatureServe 2014). Population size and trend were not used as variables in our models for similar reasons. All the detailed underpinning data for Red List evaluation are available on the BirdLife Datazone (BirdLife International 2014). Because generation time was significantly positively correlated with body size (Table S1A), we calculated and used the residual values from the simple linear regression of generation time versus body size and referred to this variable as ‘residual generation time’ following Owens and Bennett (2000).

To assess socioeconomic and demographic attributes of the countries in the parrot species’ range, we used the World Economic Outlook Database (IMF 2013), the World Factbook (CIA 2013), and the database of the Food and Agriculture Organization of the United Nations (FAOSTAT 2013). We calculated the mean values of each parameter for all countries in which a species occurred (excluding vagrant records). Industrial production growth rate, unemployment rate, human population density, urban population, human population growth rate, and agriculture area were all significantly correlated with per capita gross domestic product (GDP) (Table S1B). We used residual values for these variables after regressing them against GDP.

We analyzed the traits of 16 extinct species separately comparing trends with values for extant species.

Statistical modeling procedures

We used linear logistic regression models to identify the broad covariates of whether or not a parrot species is classified as threatened. ‘Threatened’ was defined as belonging to the IUCN threat categories ‘Vulnerable’, ‘Endangered’, and ‘Critically Endangered’, whereas ‘Non-threatened’ included the categories ‘Least Concern’ and ‘Near Threatened’. We then used ordinal regression models to evaluate in more detail the covariates of the degree of threat faced by the threatened parrot species. Both logistic and ordinal regression models were initially computed separately using each of four categories of variables (Table 1), and a final universal model was computed by combining the variables found to be significant in each of the sub-models. We used GenStat 13.7 (Payne 2009) for all modeling.

To control for phylogenetic relatedness among species we downloaded 1000 possible phylogenetic trees of *Psittaciformes* from birdtree.org. These were randomly selected to account for the branch lengths in addition to nodes separating species (Jetz et al. 2012). For each phylogeny, we ran a phylogenetic generalized least squares (PGLS) regression using the ‘caper’ package of R statistics (Freckleton et al. 2002). The explanatory and response variables were those from the best models per category found in the regression models. For each predictor in each model we report the modified p value accounting for phylogenetic relatedness, and we report λ (lambda transformation that improves the fit of the model to the phylogeny data) for each model. Greater values of λ indicate that the relationship between response and predictor variables is dependent on the phylogeny and trait values are more similar for closely related species. Values of λ closer to 0 indicate that the relationship is unrelated to phylogeny. Since these analyses were repeated for 1000 phylogenetic hypotheses, we report the mean and standard deviation of λ in each case. Additionally, we calculated in R the Blomberg’s K for threat status, which is a measure of phylogenetic signal. When values are closer to zero it indicates that the distribution of the trait is not related to phylogenetic relatedness.

We analyzed the specific threats associated with high extinction risk in parrots by assessing how many parrot species are affected by each threat type globally and specifically at the regional scale in the Neotropics, Afrotropics, Indomalaya, and Australasia/Oceania. We used threat impact scores estimated from timing, scope and severity for each threat for each species, and we compared the high and medium impact threats worldwide. We also considered the most important conservation actions needed to improve the status of threatened parrots. A formal prioritization analysis has not been carried out for all species, but the most urgent actions are documented by BirdLife International and therefore useful to present a broad level analysis.

We draw attention to countries with high risk of parrot conservation using two different methods. (1) We created a composite species ranking based on the sum of the ranks of the number of parrot species, number of globally threatened parrot species, and the number of endemic parrot species. We refer to these as ‘country priority’ scores. (2) We calculated the residual values of the final combined linear logistic regression model accounting for all significant extrinsic and intrinsic variables. We refer to these values as ‘unexplained extinction risk’ hereafter.

Results

Red list index for parrots and geographical differences

The Red List Index for parrots (0.825 in 2012) is lower than that for comparable groups of bird species including waterbirds (0.879), raptors (0.877), pigeons (0.868), gamebirds (0.842), and seabirds (0.828; Fig. 1). However, rates of decline from 1988 to 2012 are broadly comparable across these different groups: 1.27 % for parrots, 1.42 % for




Fig. 1 Red List Indices of species survival for parrots (N = 398 non data deficient extant species; *Psittaciformes*), seabirds (N = 355; *Anseriformes*, *Podicipediformes*, *Phaethontiformes*, *Gaviiformes*, *Sphenisciformes*, *Procellariiformes*, *Pelecaniformes*, *Suliformes*, *Charadriiformes*), gamebirds (N = 307; *Galliformes*), pigeons (N = 350; *Columbiformes*), raptors (N = 320; *Accipitriformes*, *Cathartiformes*, *Falconiformes*), and waterbirds (N = 852; *Anseriformes*, *Podicipediformes*, *Phoenicopteriformes*, *Gruiiformes*, *Gaviiformes*, *Ciconiiformes*, *Pelecaniformes*, *Suliformes*, *Charadriiformes*)

waterbirds, 1.75 % for raptors, 0.52 % for pigeons, 1.45 % for gamebirds, and 1.41 % for seabirds (Fig. 1).

A total of 16 parrot species are considered to be Extinct, 17 are Critically Endangered (one species is Possibly Extinct in the Wild, and another is Possibly Extinct), 40 Endangered, 54 Vulnerable, 60 Near Threatened, and 227 are categorized as Least Concern. The list of threatened parrots species with their country of occurrence and the extinct parrots are shown in the supplementary Excel data file. Species density maps of all extant parrot species (Fig S1a) and of the threatened species (Fig S1b) are shown in the online supplementary material. The Neotropics have the highest proportion of threatened species (37 % of the region's parrot species, $N = 176$), followed by Australasia/Oceania (22 %, $N = 168$ spp.), Afrotropics (19 %, $N = 26$ spp.), and Indomalaya (14 %, $N = 28$ spp.; Figure 2). Overall, there was no significant phylogenetic signal in degree of threat ($K = 0.05$, $P = 0.36$), which means that closely related species are not more likely to share the same Red List category (Fig. 3).

Threatened versus non-threatened

The probability of a parrot species being threatened decreased significantly with larger historical distribution size (combined linear logistic regression model, $\chi^2_1 = 64.89$, $P < 0.001$; Fig S2a). The probability of being threatened increased significantly with: body size ($\chi^2_1 = 13.18$, $P < 0.001$; Fig S2b), residual generation time ($\chi^2_1 = 16.82$, $P < 0.001$; Fig S2c), extent of forest dependency ($\chi^2_3 = 29.47$, $P < 0.001$; Fig S2d), and the percentage of the human population in the countries of occurrence living in urban areas ($\chi^2_1 = 28.71$, $P < 0.001$; Fig S2e). There were no significant interactions among the 27 tested variables (all $P \geq 0.184$). Table S3 lists additional variables found to be significant when analyzed separately by group (A–D). All significant variables in the combined linear logistic regression model were also significant in the PGLS model controlling for phylogeny (Table S5a).




Fig. 2 Proportion of parrot species in each IUCN Red List category for each region (following Croxall et al. 2012). Number of species in each category is shown




Fig. 3 Phylogeny of parrots indicating IUCN Red List status of each species. Colors at the tip of the branches represent the IUCN Red List category of each species; grey shading inside the circle represent major genera and groups of related genera as labeled outside the circle. Images are taken with permission from the Handbook of Birds of the World online (Taylor 1996) and represent examples of species that are critically endangered. Source of phylogeny: www.birdtree.org

Degree of threat

The severity of threat among threatened parrot species increased significantly with higher per capita GDP (Table 1) of the countries of occurrence (combined ordinal regression model, $\chi^2_1 = 8.47$, $P = 0.004$; Fig S3a) and when the species is endemic to a single country ($\chi^2_1 = 5.6$, $P = 0.018$; Fig S3b). The severity of threat was significantly less for species that are used as pets ($\chi^2_1 = 4.75$, $P = 0.029$; Fig S3c). There were no significant interactions among the 27 tested variables (all $P \geq 0.140$). Table S4 lists additional variables found to be significant when analyzed separately by group (A–D). All significant variables in the combined ordinal regression model were also significant in the PGLS model controlling for phylogeny (Table S5b).

Traits of extinct species

Sixteen parrot species in 12 genera are recorded as extinct. Five of the extinct species lived in the Neotropics (four in the Caribbean), six on islands near Africa (Mascarene Islands,

Mauritius, Seychelles), and five in Australasia/Oceania (4 in Oceania and 1 in Australia). Fourteen out of 16 extinct species (88 %) inhabited islands only. By comparison one quarter of all extant parrot species (96 out of 398 species, 24 %) inhabit only islands a significant over-representation of island species amongst those now extinct ($\chi^2_1 = 28.5$, $P < 0.001$). All but one extinct species were endemic to a single country, but distribution size was highly variable (38–557,670 km²) with the Carolina Parakeet (*Conuropsis carolinensis*) forming an outlier with range size of 3,167,000 km² (mean of 240,510 km² $\pm$ 198,000 SE for all 16 species). The mean of the extinct species' median latitude of occurrence was -6.96 degrees $\pm$ 5.00 SE compared with -6.6 degrees $\pm$ 0.75




Fig. 4 Threats: **a** percentage of parrot species impacted by the 10 major threat types. Impact is calculated from scores for timing, scope and severity (see <http://www.birdlife.org/datazone/info/spcthreat>); **b** percentage of species in each region impacted by each threat type. Only threats impacting >15 species were plotted

SE for extant species, with no significant difference (ANOVA $F_{1,413} = 0.01$, $P = 0.936$). Extinct species were mainly large bodied with a mean length of $39 \text{ cm} \pm 3.5 \text{ SE}$ (ranging between 25 and 70 cm) compared with a mean of $28.5 \text{ cm} \pm 0.7 \text{ SE}$ (ranging between 8 and 100 cm) for extant parrots (ANOVA $F_{1,410} = 7.62$, $P = 0.006$). Their estimated mean generation time was $8 \text{ years} \pm 0.7 \text{ SE}$ compared with $7.3 \text{ years} \pm 0.2 \text{ SE}$ for extant species (ANOVA $F_{1,412} = 0.92$, $P = 0.339$). All but one of the extinct species lived in only one habitat type, mainly in forests. Out of the 16 extinct species 38 % (6 species) were recorded as used as pets compared to 93 % today (372/398 species). A percentage of 69 % (11/16 spp.) of the extinct species were used as food compared to 23 % today (90/398 spp.).

Threat types, conservation actions, and priority countries

Based on the threat impact scores, the greatest threat to parrots worldwide is from agriculture (impacting 35 % of extant species), followed by hunting & trapping, logging, climate change & severe weather, and invasive alien species (Fig. 4a). The top three threats (agriculture, hunting & trapping, logging) appear roughly similar across regions, however residential development features as a greater threat in the Neotropics and Indomalaya, and invasive alien species is a greater threat in the Afrotropics and Australasia/Oceania (Fig. 4b). Whether species are threatened depends significantly on five major types of threat and their interactions: invasive alien species, agriculture, hunting & trapping, energy production & mining, and residential & commercial development (Table S6a). Higher categories of threat (i.e. Endangered, Critically Endangered) were




Fig. 5 Most important conservation actions needed by region. The figure shows for each region the percentage of species for which each conservation action is considered required by BirdLife International (2014). Only conservation actions identified as required for >15 species across all regions were plotted. Order is given by total species showing the globally most important conservation actions first

especially associated with invasive alien species, even after controlling for phylogeny ($\chi^2_1 = 14.41$, $P < 0.001$; Fig S3d, Table S6b).

The most important conservation actions as determined by BirdLife International (2014) are presented in Fig. 5. The most common actions needed in the Neotropics are site protection and management, in the Afrotropics they are legislation and ex-situ conservation, and in both Indomalaya and Australasia/Oceania they are awareness & communications, broad-scale habitat protection, and site protection (Fig. 5).

Our country priority method revealed the following 10 countries with highest priority ranks: Indonesia, Brazil, Australia, Colombia, Bolivia, Ecuador, Peru, Papua New Guinea, Venezuela, and Mexico (more countries in Table 2). Examination of the unexplained extinction risk (after all significant intrinsic and extrinsic variables were accounted for in our models—see above) showed that island countries (or territories) accounted for both the lowest and highest values (Fig. 6). In total 40 countries had more threatened species than predicted (i.e. positive residuals), 17 of these were islands, 21 were in Africa. The Philippines (Indomalaya) and New Zealand (Australasia) also showed positive residuals (Fig. 6). In the Americas, all countries had less threatened species than predicted (with Brazil being closest to zero; Fig. 6).

Table 2 Countries of highest priority for parrot conservation, ranked by total numbers of (a) parrot species, (b) globally threatened species, and (c) endemic species (restricted to one country)

Country	Diversity rank	Threat rank	Endemics rank	Overall rank
Indonesia	1	2	2	1
Brazil	2	1	3	2
Australia	4	6	1	3.5
Colombia	3	2	6	3.5
Bolivia	5	4	8	5
Ecuador	8	3	9	6.5
Peru	6	5	9	6.5
Papua New Guinea	6	11	5	8
Venezuela	7	8	8	9
Mexico	13	6	8	10
Philippines	16	8	4	11
Argentina	10	9	11	12
Guyana	9	11	11	13.5
Panama	11	10	10	13.5
New Zealand	20	7	5	15.5
Paraguay	12	9	11	15.5
French Guiana	10	13	11	17.5
Suriname	10	13	11	17.5
Costa Rica	14	10	11	19
Guatemala	15	11	11	20

Overall rank is derived from the sum of ranks for the three parameters




Fig. 6 Unexplained (residual) extinction risk of parrots by country and region: values shown are each country's mean residuals from the combined linear logistic regression model predicting likelihood of a parrot species being threatened (VU, EN, CR). High positive values indicate high unexplained extinction risk

Discussion

Our analysis of Red List Indices revealed that parrots are more threatened than other comparable taxonomic groups (Fig. 1) with consistently negative trends in extinction risk over the last 25 years. The results show that conservation successes have been outweighed by the number of species being up-listed to higher categories of threat (IUCN 2014). Importantly, our analysis showed no significant effect of phylogeny on threatened status of parrots (Fig. 3). Extinction risk is usually non-randomly distributed with respect to phylogeny (Fisher and Owens 2004), hence the most illuminating comparative models are those like ours that focus on relatively narrow taxonomic groups (Cardillo et al. 2008).

Biological attributes

We found several biological attributes associated with extinction risk in parrots. We found that species with larger historical distributions are less likely to be threatened (Fig S2a). This probably reflects the link between historical and present distribution size, and the fact that widely distributed species are often adapted to multiple habitats and less impacted by local threats (Ewers and Didham 2006; Purvis et al. 2000).

As for other taxonomic groups, large bodied parrots are more prone to extinction risk (Fig S2b) (Bennett and Owens 1997; Cardillo 2003; Cardillo and Bromham 2001; Purvis et al. 2000). Large body size correlates with many known extinction-promoting traits, including low population densities, slower life histories (Cardillo et al. 2005), and higher risk from hunters (Cowlshaw and Dunbar 2000). The significant effect of residual generation time also indicates that parrots with slower life history (longer generation time) are more likely to be threatened, independently of body size (Fig S2c). Bennett and Owens (1997) also found that increasing body size and residual generation time were associated with increased extinction risk among birds when analyzed at the family level. Marsden and Royle (2015) showed that larger bodied parrots are predictably uncommon in the wild.

Our study confirmed that forest dependent parrot species are more likely to be threatened (Fig S2d). Most parrots are forest dependent because of their nesting and feeding habits (Snyder et al. 2000; White et al. 2005). At least 70 % of parrots are secondary tree cavity nesters (nesting in preexisting tree cavities), hence primary forest destruction decreases the number of available nest sites and reproductive success (Newton 1994). Woodpeckers, as potentially important keystone cavity excavators in Neotropical forests (Cornelius et al. 2008), can also be threatened by anthropogenic factors that ultimately affect parrot reproduction as well. Forest destruction can also lead to the loss of key food resources, as many parrots eat mostly tree seeds and fruits and rely on large areas of suitable habitat to provide year-round sources of these temporally variable food sources (Arndt 2007; Brightsmith 2005; Forshaw 2011; Juniper and Parr 2003).

Anthropogenic and socio-economic factors

Our data suggest a subtle effect of the extent of economic development of the countries where the parrots live. Whereas the extent of urbanization (percentage of human populations living in urban areas) seems to relate broadly to whether a parrot species is threatened (Fig S2e), the impact of per capita GDP (i.e. developed economies) seems to have most influence by pushing parrot species into higher categories of extinction risk (Fig S3a). Urbanization can entail far-reaching transformations in itself but is also linked to broad scale environmental degradation (Cohen 2006; McKinney 2002). As populations and economies of primarily rural countries grow, there is often a migration of people to urban areas, and this in turn can be followed by further economic expansion and higher per capita GDP (Chang and Brada 2006; Moomaw and Shatter 1996). Hence urbanization can occur at high levels before GDP is maximized (Brühlhart and Sbergami 2009; Henderson 2003). Our results suggest that urbanization is a good variable accounting for the impact on parrots of the earlier stages of economic development, with the continuing impact of economic development best captured by GDP.

Parrots with distributions limited to one country are also more likely to belong to higher categories of endangerment (Fig S3b). Our analysis thus suggests that extinction risk of single country endemics (45 % of all parrot species) is often adversely affected by either

the singular history or the conservation management practices (or lack thereof) of their single country of occurrence. In the latter case, specific threats may be better controlled in some nations than others (e.g. progressive forestry laws and control of pet trade) with potential advantages to species living in more than one country (Hirakuri 2003; Pires 2012; Ribot 1999; Sunderlin et al. 2005).

Our finding that threatened species used for pets tend to belong to the lower categories of endangerment may seem counterintuitive (Fig S3c) but is supported by recent studies which show that the vast majority of species in domestic and international bird trade are non-threatened species (Pires and Clarke 2012; Pires 2012) and that utilized species in general are less threatened (Butchart 2008). This finding likely speaks much more to how increased threatened status (or decreased abundance) reduces use as pets than it does to how use for pets causes increases in threatened status. Pires and Clarke (2012) report that rare parrot species are only infrequently found in illegal parrot markets, confirming that most parrot poaching is driven by species that are easier to catch rather than by commercially based poachers searching for the rarest species.

Traits of extinct species

Our analysis of the traits of extinct species provides additional insight on some of our key results concerning extinction risk of extant species. For example, a disproportionate number of extinct parrot species (88 % vs. 24 % for extant species) were insular, yet our models for extant species did not show that island endemism was a significant factor. This suggests that island endemism has been a strong factor in extinction risk but may have been masked in our analyses of extant species because many of the most susceptible species have already become extinct. In keeping with our models, extinct parrot species were larger on average than extant parrots, and most extinct species (94 %) were single country endemics (compared to 45 % for extant species).

Threat types and conservation actions

Our analysis reveals that the most important threats to parrots are agriculture, hunting and trapping, logging, climate change and severe weather, invasive alien species, and residential and commercial development (Fig. 4a, Table S6a). Invasions by non-native species are often cited as leading causes of species extinctions (Carrete and Tella 2008; Gurevitch and Padilla 2004), but our analysis suggests more complex synergistic effects in combination with other threats, including hunting and trapping, and agriculture (see interactions in Table S6a). Interestingly, ‘invasive alien species’ was the sole threat significantly associated with the higher categories of threat status (Fig S3d, Table S6b). This warrants increasing conservation attention on invasive alien species. These threats do not seem to respond to the major conservation strategies that are normally applied (e.g. country border protection, legislation, law enforcement). Rather invasive species are a threat that needs to be combatted directly and individually through regionally designed conservation action plans.

We found regional differences in the threats impacting parrots and hence the identified conservation actions required. Agriculture is a particularly important threat worldwide in terms of number of parrot species impacted (Fig. 4a), and especially in the Neotropics (Fig. 4b) where site protection is identified as the major conservation action required (Fig. 5). In Indomalaya and Australasia/Oceania, the most significant threat to parrots is logging (Fig. 4b), driving extensive deforestation in recent decades (Sodhi et al. 2004,

2010a, b). Reflecting this, the most common conservation actions needed are broad-scale habitat protection, site protection and awareness raising (Fig. 5). In contrast, the most common threat in the Afrotropics is hunting and trapping (primarily for the cage-bird trade and for use as pets; Fig. 4b), and the most important key actions include enforcing and enhancing legislation (Fig. 5).

Priority countries

Countries with the highest conservation priority were from all regions except the Afrotropics. Although Indonesia ranked the highest overall and Australia had the highest value amongst developed countries, 15 out of the top 20 parrot conservation priority countries were from the Neotropics (Table 2). These countries deserve high conservation interest given the high diversity and endemism of their parrot species, and the high proportion of their parrot species that are threatened. After accounting for all the significant variables in our model (historical distribution size, body size, generation time, forest dependency, and percentage of urban human population in the distribution ranges), the mean residual (or unexplained) extinction risk values of each country suggested some variation in country-level performances. The countries with positive residuals (on average more threatened species than expected) included many islands of different sizes (island nations, Philippines, and New Zealand; Fig. 6). Interestingly, island countries and territories also had the lowest mean residuals (less threatened species than expected; e.g. Bahamas, Comoros, Micronesia) in conservation status (Fig. 6). Considering only mainland countries, Africa showed the worst trends compared to the other regions, where capture for the pet trade is still common (Fig. 6). These results again confirm the impact on parrots of living on islands, and our method may prove important for identifying poorly performing countries (e.g. Dominican Republic, Haiti, Brazil, Burundi, Philippines, New Zealand, etc.) that need both extra conservation attention and further research on the nature of threatening processes.

Conclusion

Our study confirms that parrots are more threatened on average than comparable bird species groups, and that biological factors known to affect extinction risk in other taxa also apply to parrots. Thus parrot species with small historical distribution size, large body size, long generation time, and forest dependency are most likely to be categorized as threatened (Vulnerable, Endangered, Critically Endangered) under IUCN Red List criteria. Extinct parrot species shared most of these traits but also highlight island endemism as an important factor in the past. The extent of a country's urbanization provides a broad proxy for the major human socio-economic drivers of extinction risk in parrots. Our models also revealed that parrots are more likely to be highly threatened if their distribution falls within a single country's jurisdiction and in countries with higher per capita GDP, presumably because the higher levels of development that these factors are associated with, tend to drive the major threats to parrots worldwide including agriculture, hunting and trapping, and logging.

Acknowledgments The authors thank the many thousands of individuals and organizations who contribute to BirdLife International's assessments of parrots for the IUCN Red List, upon which the analyses presented here are based. We thank Hannah Wheatley for the management of these data, Mark Balam and

Janet Nackoney for help with GIS mapping, Rod Peakall for commenting on the manuscript, Kurt Holle and Jenő Gászpor for helping with the socio-economic variables, and Marta Romhányi, Attila Romhányi, and Braulio Poje for helping with the database construction.

References

- Arndt T (2007) *Lexicon of Parrots*. Arndt Verlag, Bretten
- Baker-Gabb D (2011) National Recovery Plan for the Superb Parrot *Polytelis swainsonii*. Department of Sustainability and Environment, Melbourne
- Beissinger SR, Bucher EH (1992) Can parrots be conserved through sustainable harvesting? *Bioscience* 42:164–173
- Beissinger SR, Snyder NFR, Thomsen JB, Mulliken TA (1992) Trade in neotropical psittacines and its conservation implications. *New world parrots in crisis: solutions from conservation biology*. Smithsonian Institution Press, New York and London, pp 221–239
- Bennett PM, Owens IPF (1997) Variation in extinction risk among birds: chance or evolutionary predisposition? *Proc R Soc Lond B Biol Sci* 264:401–408. doi:10.1098/rspb.1997.0057
- BirdLife International (2014) The BirdLife checklist of the birds of the world: Version 7. http://www.birdlife.org/datazone/userfiles/file/Species/Taxonomy/BirdLife_Checklist_Version_70.zip
- BirdLife International and NatureServe (2014) Bird species distribution maps of the world. Version 4.0. BirdLife International, Cambridge, UK and NatureServe, Arlington, USA. <http://www.birdlife.org/datazone/info/spcdownload>
- Blaikie P, Jeanrenaud S (1997) Biodiversity and human welfare. *Social change and conservation*. Earthscan, London, pp 46–70
- Brightsmith DJ (2005) Competition, predation and nest niche shifts among tropical cavity nesters: ecological evidence. *J Avian Biol* 36:74–83
- Brühlhart M, Sbergami F (2009) Agglomeration and growth: cross-country evidence. *J Urban Econ* 65:48–63. doi:10.1016/j.jue.2008.08.003
- Bush ER, Baker SE, Macdonald DW (2014) Global trade in exotic pets 2006–2012. *Conserv Biol* 28:663–676. doi:10.1111/cobi.12240
- Butchart SHM (2008) Red List Indices to measure the sustainability of species use and impacts of invasive alien species. *Bird Conserv Int* 18:S245–S262. doi:10.1017/S095927090800035X
- Butchart SHM et al (2004) Measuring global trends in the status of biodiversity: Red List Indices for birds. *PLoS Biol* 2:383
- Butchart SHM et al (2005) Using Red List Indices to measure progress towards the 2010 target and beyond. *Philos Trans R Soc B* 360:255–268. doi:10.1098/rstb.2004.1583
- Butchart SHM et al (2007) Improvements to the Red List Index. *PLoS One* 2:e140. doi:10.1371/journal.pone.0000140
- Butchart SHM et al (2010) Global biodiversity: indicators of recent declines. *Science* 328:1164–1168. doi:10.1126/science.1187512
- Cardillo M (2003) Biological determinants of extinction risk: why are smaller species less vulnerable? *Anim Conserv* 6:63–69
- Cardillo M, Bromham L (2001) Body size and risk of extinction in Australian mammals. *Conserv Biol* 15:1435–1440
- Cardillo M, Purvis A, Sechrest W, Gittleman JL, Bielby J, Mace GM (2004) Human population density and extinction risk in the world's carnivores. *PLoS Biol* 2:e197. doi:10.1371/journal.pbio.0020197
- Cardillo M et al (2005) Multiple causes of high extinction risk in large mammal species. *Science* 309:1239–1241. doi:10.1126/science.1116030
- Cardillo M, Mace GM, Gittleman JL, Jones KE, Bielby J, Purvis A (2008) The predictability of extinction: biological and external correlates of decline in mammals. *Proc R Soc B* 275:1441–1448. doi:10.1098/rspb.2008.0179
- Carrete M, Tella JL (2008) Wild-bird trade and exotic invasions: a new link of conservation concern? *Front Ecol Environ* 6:207–211
- Chang GH, Brada JC (2006) The paradox of China's growing under-urbanization. *Econ Syst* 30:24–40. doi:10.1016/j.ecosys.2005.07.002
- CIA (2013) Central Intelligence Agency: The World Factbook. <https://www.cia.gov/library/publications/the-world-factbook/>. Accessed on June 2013
- Cohen B (2006) Urbanization in developing countries: Current trends, future projections, and key challenges for sustainability. *Technol Soc* 28:63–80. doi:10.1016/j.techsoc.2005.10.005

- Collar NJ (1997) Family Psittacidae (parrots). In: Hoyo Jd, Elliott A, Sargatal J (eds) Handbook of the birds of the world, vol 4. Lynx Edicions, Barcelona, pp 280–477
- Collar NJ (2000) Globally threatened parrots: criteria, characteristics and cures. *Int Zoo Yearb* 37:21–35
- Collar NJ, Juniper AT (1992) Dimensions and causes of the parrot conservation crisis. In: Beissinger SR, Snyder NFR (eds) *New World parrots in crisis: solutions from conservation biology*. Smithsonian Institution Press, Washington, DC, pp 1–24
- Cornelius C et al (2008) Cavity-nesting birds in Neotropical forests: cavities as a potentially limiting resource. *Ornitol Neotrop* 19:253–268
- Cowlishaw G, Dunbar RIM (2000) *Primate conservation biology*. University of Chicago Press, Chicago
- Croxall JP, Butchart SHM, Lascelles B, Stattersfield AJ, Sullivan B, Symes A, Taylor P (2012) Seabird conservation status, threats and priority actions: a global assessment. *Bird Conserv Int* 22:1–34. doi:[10.1017/S0959270912000020](https://doi.org/10.1017/S0959270912000020)
- Ewers RM, Didham RK (2006) Confounding factors in the detection of species responses to habitat fragmentation. *Biol Rev* 81:117–142. doi:[10.1017/S1464793105006949](https://doi.org/10.1017/S1464793105006949)
- FAOSTAT (2013) Food and Agriculture Organization of the United Nations. <http://faostat3.fao.org/>. Accessed on July 2013
- Fisher DO, Owens IPF (2004) The comparative method in conservation biology. *Trends Ecol Evol* 19:391–398. doi:[10.1016/j.tree.2004.05.004](https://doi.org/10.1016/j.tree.2004.05.004)
- Fisher DO, Blomberg SP, Owens IPF (2003) Extrinsic versus intrinsic factors in the decline and extinction of Australian marsupials. *Proc R Soc Lond B* 270:1801–1808. doi:[10.1098/rspb.2003.2447](https://doi.org/10.1098/rspb.2003.2447)
- Forshaw JM (2011) *Parrots of the world* CSIRO Publishing, Collingwood, Australia
- Freckleton RP, Harvey PH, Pagel M (2002) Phylogenetic analysis and comparative data: a test and review of evidence. *Am Nat* 160:712–726. doi:[10.1086/343873](https://doi.org/10.1086/343873)
- Grahl Wd (1990) *Papageien: Lebensweise, Arten, Zucht*, 9th edn. Verlag Eugen Ulmer, Germany
- Gurevitch J, Padilla DK (2004) Are invasive species a major cause of extinctions? *Trends Ecol Evol* 19:470–474. doi:[10.1016/j.tree.2004.07.005](https://doi.org/10.1016/j.tree.2004.07.005)
- Harris JB et al (2012) Managing the long-term persistence of a rare cockatoo under climate change. *J Appl Ecol* 49:785–794
- Henderson V (2003) The urbanization process and economic growth: the so-what question. *J Econ Growth* 8:47–71. doi:[10.1023/A:1022860800744](https://doi.org/10.1023/A:1022860800744)
- Hirakuri SR (2003) *Can law save the forest?: lessons from Finland and Brazil*. CIFOR, Bogor
- Holdsworth M, Starks J (2006) *National Recovery Plan for the Orange-bellied Parrot (Neophema chrysogaster)*. Department of Primary Industries and Water (DPIW), Hobart, Australia
- IMF (2013) International Monetary Fund: world economic outlook database. <http://www.imf.org/>. Accessed on June 2013
- IUCN (2014) The IUCN Red list of threatened species. Version 2014.2. <http://www.iucnredlist.org>
- Jetz W, Thomas GH, Joy JB, Hartmann K, Mooers AO (2012) The global diversity of birds in space and time. *Nature* 491: 444–448. <http://www.nature.com/nature/journal/v491/n7424/abs/nature11631.html> - supplementary-information
- Jones M, Fielding A, Sullivan M (2006) Analysing extinction risk in parrots using decision trees. *Biodivers Conserv* 15:1993–2007. doi:[10.1007/s10531-005-4316-1](https://doi.org/10.1007/s10531-005-4316-1)
- Juniper T, Parr M (2003) *Parrots: a guide to parrots of the world*. A & C Black, USA
- Kerr JT, Currie DJ (1995) Effects of human activity on global extinction risk. *Conserv Biol* 9:1528–1538. doi:[10.1046/j.1523-1739.1995.09061528.x](https://doi.org/10.1046/j.1523-1739.1995.09061528.x)
- Laurance WF, Albernaz AKM, Schroth G, Fearnside PM, Bergen S, Venticinque EM, Da Costa C (2002) Predictors of deforestation in the Brazilian Amazon. *J Biogeogr* 29:737–748. doi:[10.1046/j.1365-2699.2002.00721.x](https://doi.org/10.1046/j.1365-2699.2002.00721.x)
- Laurance WF, Goosem M, Laurance SGW (2009) Impacts of roads and linear clearings on tropical forests. *Trends Ecol Evol* 24:659–669. doi:[10.1016/j.tree.2009.06.009](https://doi.org/10.1016/j.tree.2009.06.009)
- Marsden SJ, Royle K (2015) Abundance and abundance change in the world's parrots. *Ibis* 157:219–229. doi:[10.1111/ibi.12236](https://doi.org/10.1111/ibi.12236)
- Martin RO et al (2014) Research and conservation of the larger parrots of Africa and Madagascar: a review of knowledge gaps and opportunities. *Ostrich* 85:205–233. doi:[10.2989/00306525.2014.948943](https://doi.org/10.2989/00306525.2014.948943)
- McKinney ML (2002) Urbanization, biodiversity, and conservation: the impacts of urbanization on native species are poorly studied, but educating a highly urbanized human population about these impacts can greatly improve species conservation in all ecosystems. *Bioscience* 52:883–890. doi:[10.1641/0006-3568\(2002\)052\[0883:ubac\]2.0.co;2](https://doi.org/10.1641/0006-3568(2002)052[0883:ubac]2.0.co;2)
- Moomaw RL, Shatter AM (1996) Urbanization and economic development: a Bias toward Large Cities? *J Urban Econ* 40:13–37. doi:[10.1006/juec.1996.0021](https://doi.org/10.1006/juec.1996.0021)

- Müller M (2000) Review of the in situ status of the Great green or Buffon's macaw *Ara ambigua* and the European Endangered Species Programme (EEP). *Int Zoo Yearb* 37:183–190
- Newton I (1994) The role of nest sites in limiting the numbers of hole-nesting birds: a review. *Biol Conserv* 70:265–276. doi:[10.1016/0006-3207\(94\)90172-4](https://doi.org/10.1016/0006-3207(94)90172-4)
- Olson DM et al (2001) Terrestrial ecoregions of the world: a new map of life on earth. *Bioscience* 51:933–938. doi:[10.1641/0006-3568\(2001\)051\[0933:TEOTWA\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2001)051[0933:TEOTWA]2.0.CO;2)
- Owens IPF, Bennett PM (2000) Ecological basis of extinction risk in birds: habitat loss versus human persecution and introduced predators. *Proc Natl Acad Sci* 97:12144–12148. doi:[10.1073/pnas.200223397](https://doi.org/10.1073/pnas.200223397)
- Pain DJ et al (2006) Impact of protection on nest take and nesting success of parrots in Africa, Asia and Australasia. *Anim Conserv* 9:322–330. doi:[10.1111/j.1469-1795.2006.00040.x](https://doi.org/10.1111/j.1469-1795.2006.00040.x)
- Pasquier RF (1980) Conservation of new world parrots: proceedings of the ICBP Parrot Working Group Meeting, St. Lucia, 1980. Smithsonian Institution Press for the International Council for Bird Preservation, USA
- Payne RW (2009) *GenStat*. Wiley Interdisciplinary Reviews: Computational Statistics 1:255–258. doi:[10.1002/wics.32](https://doi.org/10.1002/wics.32)
- Pimm SL et al (2014) The biodiversity of species and their rates of extinction, distribution, and protection. *Science*. doi:[10.1126/science.1246752](https://doi.org/10.1126/science.1246752)
- Pires SF (2012) The illegal parrot trade: a literature review. *Glob Crime* 13:176–190
- Pires S, Clarke RV (2012) Are parrots CRAVED? An analysis of parrot poaching in Mexico. *J Res Crime Delinq* 49:122–146
- Purvis A, Gittleman JL, Cowlishaw G, Mace GM (2000) Predicting extinction risk in declining species. *Proc R Soc Lond B Biol Sci* 267:1947–1952. doi:[10.1098/rspb.2000.1234](https://doi.org/10.1098/rspb.2000.1234)
- Ribot JC (1999) Decentralisation, participation and accountability in Sahelian forestry: legal instruments of political-administrative control. *Africa* 69:23–65. doi:[10.2307/1161076](https://doi.org/10.2307/1161076)
- Snyder N, McGowan P, Gilardi J, Grajal A (2000) Parrots. Status Survey and Conservation Action Plan 2000–2004. IUCN, Gland, Switzerland and Cambridge, UK
- Sodhi NS, Koh LP, Brook BW, Ng PKL (2004) Southeast Asian biodiversity: an impending disaster. *Trends Ecol Evol* 19:654–660. doi:[10.1016/j.tree.2004.09.006](https://doi.org/10.1016/j.tree.2004.09.006)
- Sodhi N, Posa M, Lee T, Bickford D, Koh L, Brook B (2010a) The state and conservation of Southeast Asian biodiversity. *Biodivers Conserv* 19:317–328. doi:[10.1007/s10531-009-9607-5](https://doi.org/10.1007/s10531-009-9607-5)
- Sodhi NS et al (2010b) Deforestation and avian extinction on tropical landbridge islands. *Conserv Biol* 24:1290–1298. doi:[10.1111/j.1523-1739.2010.01495.x](https://doi.org/10.1111/j.1523-1739.2010.01495.x)
- Spray SL, McGlothlin KL (2003) Loss of biodiversity. Exploring environmental challenges. Rowman & Littlefield, Lanham
- Sunderlin WD, Angelsen A, Belcher B, Burgers P, Nasi R, Santoso L, Wunder S (2005) Livelihoods, forests, and conservation in developing countries: an overview. *World Dev* 33:1383–1402. doi:[10.1016/j.worlddev.2004.10.004](https://doi.org/10.1016/j.worlddev.2004.10.004)
- Taylor B (1996) Sulu Racquet-tail (*Prioniturus verticalis*), White Cockatoo (*Cacatua alba*), New Zealand Kaka (*Nestor meridionalis*), Blue-throated Macaw (*Ara glaucogularis*), Grey-breasted Parakeet (*Pyrrhura griseipectus*), Puerto Rican Amazon (*Amazona vittata*), Orange-bellied Parrot (*Neophema chrysogaster*), Malherbe's Parakeet (*Cyanoramphus malherbi*), Red-throated Lorikeet (*Charmosyna amabilis*). In: del Hoyo J, Elliott A, Sargatal J, Christie DA, de Juana E (eds) (2013). Handbook of the Birds of the World Alive. Lynx Edicions, Barcelona. <http://www.hbw.com>. Accessed on 12 Nov 2014
- Tella JL, Hiraldo F (2014) Illegal and legal parrot trade shows a long-term, cross-cultural preference for the most attractive species increasing their risk of extinction. *PLoS One* 9:e107546. doi:[10.1371/journal.pone.0107546](https://doi.org/10.1371/journal.pone.0107546)
- Tittensor DP et al (2014) A mid-term analysis of progress toward international biodiversity targets. *Science* 346:241–244. doi:[10.1126/science.1257484](https://doi.org/10.1126/science.1257484)
- Webster A, Price R, Lowe K (2003) Action statement: Swift Parrot (*Lathamus discolor*). Department of Sustainability and Environment, Victoria
- White TH Jr, Collazo JA, Vilella FJ (2005) Survival of captive-reared Puerto Rican parrots released in the Caribbean National Forest. *The Condor* 107:424–432

ONLINE SUPPLEMENTARY MATERIAL

Supplementary Methods

Red List Indices

We examined trends in extinction risk during 1988-2012 for parrots (*Psittaciformes*) and comparable high profile species-groups with similar numbers of species using Red List Indices. These groups included waterbirds (including species from families *Anseriformes*, *Podicipediformes*, *Phoenicopteriformes*, *Gruiformes*, *Gaviiformes*, *Ciconiiformes*, *Pelecaniformes*, *Suliformes*, *Charadriiformes*), seabirds (*Anseriformes*, *Podicipediformes*, *Phaethontiformes*, *Gaviiformes*, *Sphenisciformes*, *Procellariiformes*, *Pelecaniformes*, *Suliformes*, *Charadriiformes*), raptors (*Accipitriformes*, *Cathartiformes*, *Falconiformes*), each of which comprises multiple orders. Other groups included the largest bird orders (i.e. with more than 250 species) like pigeons (*Columbiformes*) and gamebirds (*Galliformes*), except the orders *Passeriformes*, *Caprimulgiformes*, and *Piciformes*. Cases where species were re-categorized owing to improved knowledge or revised taxonomy are excluded. We used data from the comprehensive assessments of all bird species in 1988, 1994, 2000, 2004, 2008 and 2012, updated to 2014 (Tittensor et al. 2014).

Database and variables

We assembled a database of the biological and geographic attributes of all 398 extant parrot species using the 2014 version of BirdLife International and IUCN's database which underpins the IUCN Red List assessments for birds on the BirdLife Data Zone (BirdLife International 2014) and IUCN Red List website (IUCN 2014). We added further data including the socio-economic and demographic attributes of the countries the parrots occur in, from various external sources (Table 1).

We used the IUCN Red List extinction risk categories of all extant species of parrots (BirdLife International 2014; IUCN 2014) as the response variables in our analyses. In tables and graphs we use the standard IUCN abbreviations for Red List categories as follows: LC = Least Concern, NT = Near Threatened, VU = Vulnerable, EN = Endangered, CR = Critically Endangered (IUCN 2014). We analyzed the traits of 16 extinct species separately, and exclude hypothetical taxa that have not been confirmed as valid species (Hoyo et al. 2014). For all analyses we followed the taxonomy of BirdLife International (2014).

Because of the large number of explanatory variables we initially divided the potential explanatory variables into four groups and performed analyses separately for each. We then combined all significant variables from each sub-analysis into a final model (see below). The groups were: (A) geographical and distributional attributes of each species; (B) biological, ecological and life history variables; (C) type of utilization by humans; and (D) socio-economic and demographic attributes of the countries where the species occur. For detailed descriptions, source of data and values of each variable see Table 1.

We used spatial analyses on the digital distribution files from BirdLife International and NatureServe (2014). ArcGIS 10.2 was used to calculate the median latitude of the distribution of each species. We calculated historical distribution size (i.e. current plus extirpated or historical ranges) from the species' shape-files. We used historical distribution size instead of current distribution sizes in order to avoid possible circularity as current extent of occurrence is a parameter used in the IUCN Red List criteria. For the same reason population size and trend were not used, to avoid circularity (IUCN 2014).

We defined whether each species was an island endemic (yes/no) depending on whether it was restricted to an island smaller than 110 000 km². Under this arbitrary definition, parrots of larger islands such as Borneo (743 330 km²), Sumatra (473 481 km²), or New Guinea (452 860 km²) were not considered island endemics; the largest island that qualified was Cuba

(109 820 km²) (see description in Table 1). We tested the validity of this assumption by varying our definition of the island size (including larger island cutoffs) that qualified and found this made no difference to the results.

We determined the type of utilization by people (group C) from the IUCN Red List Use Classification Scheme (<http://www.iucnredlist.org/technical-documents/classification-schemes>) assigned into binomial variables (yes/no). Pets are defined as those species recorded as being kept in captivity, either as personal pets, or for display in zoos, collections etc.

To assess socioeconomic and demographic attributes, we used The World Economic Outlook Database (IMF 2013), The World Factbook (CIA 2013), and the database of the Food and Agriculture Organization of the United Nations (FAOSTAT 2013) as sources, and calculated the mean values of each parameter for all countries in which a species occurred (excluding vagrant records; Table 1).

Historical distribution size and body size were normalized using a log_e transformation.

Threats, conservation actions, and priority countries

To understand the specific threats associated with high extinction risk in parrots, we extracted data from BirdLife International (2014) who classify threats using the IUCN-CMP Unified Classification of Direct Threats (Salafsky et al. 2009, updated at <http://www.iucnredlist.org/technical-documents/classification-schemes/threats-classification-scheme>). We analyzed threats at level 1, apart from Biological Resource Use and Natural System Modifications, which we analyzed at level 2 given the fundamentally distinct processes these classes aggregate. We assessed how many parrot species are affected by each threat type globally and at the regional scale. We also considered the overall threat impact scores (which are calculated from the timing, scope and severity of

each threat to each species:

http://www.iucnredlist.org/documents/Dec_2012_Guidance_on_Threat_Impact_Scoring.pdf

f), and excluded past and unknown threats and those with no/negligible impacts.

We analyzed data from BirdLife International (2014) on the most important conservation actions needed to improve the status of threatened parrots; these are coded against the IUCN-CMP Unified Classification of Actions (Salafsky et al. 2009, updated at http://www.iucnredlist.org/documents/Dec_2012_Guidance_Conservation_Actions_Needed_Classification_Scheme.pdf).

We used two methods to highlight important countries for parrot conservation. (1) We followed Croxall et al. (2012) to prioritize countries according to the sum of their ranks for the total numbers of their (a) parrot species, (b) globally threatened species, and (c) single country endemic species, and referred as 'country priority'. (2) In order to determine which countries had the highest proportion of unexplained extinction risk we calculated the mean for each country of the residuals from the combined linear logistic regression (see below) and used this to rank them in terms of the magnitude of unexplained variation remaining once all known significant causes of threatened status have been removed. We refer to this as 'unexplained extinction risk'.

Statistical analysis

We conducted our analyses of the likely determinants of the status of parrots at two levels, one designed to identify the broad covariates of whether a parrot species is threatened or not, and the other designed to evaluate in more detail the covariates of the degree of threat faced by parrot species. To test variables at the broader scale we assigned all species a binary response variable of 0 (Least Concern and Near Threatened) or 1 ('Threatened', i.e. Vulnerable, Endangered or Critically Endangered), and analyzed possible explanatory

variables using linear logistic regression with appropriate controls for non-independence due to phylogenetic effects (see below).

To examine the possible causes of threat in further detail, we assigned numerical values corresponding to the extinction risk faced by each threatened species as follows: Vulnerable = 1, Endangered = 2, Critically Endangered = 3. We used ordinal regression models to analyze these values because of their directional numerical nature.

Linear logistic regression and ordinal regression models were initially computed using each set of variables (A-D above) separately. This was to avoid statistical issues associated with multicollinearity. The final universal model was computed by combining the variables found to be significant in each of the sub-models.

We used correlation matrices to determine whether variables within each group were correlated, and initially avoided using correlated variables in the same analysis. The following variables were excluded on this basis: mean body mass and clutch size (correlated with body size, Table S1a) and forest area of each country (correlated with area of agriculture, Table S1b). Because generation time was significantly positively correlated with body size (Table S1a), we calculated and used the residual values from the simple linear regression of generation time versus body size and referred this variable as 'residual generation time' following Owens and Bennett (2000). Similarly, we calculated residual values for industrial production growth rate, unemployment rate, human population density, urban population, human population growth rate, and agriculture area because they were significantly correlated with per capita GDP (Table S1b).

All linear logistic regressions and ordinal regression models were computed using GenStat 13.7 (Payne 2009). Akaike information criteria (AIC) and Bayesian information criteria (BIC) were used to determine the best parsimonious models containing all significant terms. Models were selected with the lowest AIC values and simultaneously having the lowest BIC

values (Table S2). We also report *P*-values for each significant variable determined by its exclusion from the full models selected above. We also controlled for phylogenetic relatedness between species using phylogenetic generalized least squares (PGLS) regression.

Supplementary Results

Table S1. Correlation matrices. Correlation coefficients (Pearson's) are shown below diagonal, two-sided test of correlations different from zero above diagonal.

(a) Number of species: 337

Variables	Body Size	Body Mass	Clutch Size	Generation Time
Body Size	-	<0.001	<0.001	<0.001
Body Mass	0.8447	-	<0.001	<0.001
Clutch Size	-0.2214	-0.3011	-	<0.001
Generation Time	0.6533	0.7552	-0.2535	-

(b) Number of species: 378

Variables	Per capita GDP	Industrial Production Growth Rate	Unemployment Rate	Human Population Density	Urban Population	Human Population Growth Rate	Agriculture Area	Forest Area
Per capita GDP	-	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
Industrial Production Growth Rate	-0.5775	-	0.5399	0.1514	<0.001	<0.001	<0.001	<0.001
Unemployment Rate	-0.2218	0.0316	-	0.5357	0.0343	<0.001	<0.001	0.0019
Human Population Density	-0.373	0.0739	0.032	-	<0.001	0.0189	0.9509	0.0068
Urban Population	0.6685	-0.6265	-0.1089	-0.5035	-	<0.001	<0.001	<0.001
Human Population Growth Rate	-0.3382	0.4275	0.3233	0.1207	-0.55	-	0.7567	0.0039
Agriculture Area	0.4914	-0.5284	0.2471	-0.0032	0.4027	-0.016	-	<0.001
Forest Area	-0.5472	0.3489	-0.1589	-0.139	-0.2195	-0.1481	-0.8451	-

Table S2. AIC (Akaike information criterion) and BIC (Bayesian information criterion, also referred to as Schwarz information criterion) values for best parsimonious models containing all significant terms for each variable group (ranked by AIC values).

Model	Linear Logistic Regression		
	AIC	BIC	d.f.
Threats	249.23	273.15	6
A - Geographical and distributional attributes	281.64	304.29	6
Final combined model	292.42	324.29	8
B - Biological, ecological and life history variables	327.60	354.02	7
D - Socio-economic and demographic attributes	375.71	391.45	4
C - Type of utilization by humans	452.62	468.55	4

Model	Ordinal Regression		
	AIC	BIC	d.f.
B - Biological, ecological and life history variables	358.22	664.74	96
A - Geographical and distributional attributes	360.82	670.54	97
D - Socio-economic and demographic attributes	382.42	719.01	103
Final combined model	432.68	822.95	115
Threats	436.95	828.26	115
C - Type of utilization by humans	440.57	824.05	113

Table S3. Significant variables in the linear logistic regression models run separately for each variable group (A-D) predicting the likelihood of a parrot species being threatened (VU, EN, CR). λ values for the phylogenetic generalized least squares model (PGLS) are given in the table for each model. We give mean $\pm$ standard deviation (SD).

Group	Variable	Wald statistic	d.f.	$P (\chi^2)$	Estimate $\pm$ SD (PGLS)	P (PGLS) $\pm$ SD	$\lambda \pm$ SD
A	Historical Distribution Size ($\log_e$)	71.64	1	<0.001	-0.246 $\pm$ 0.004	<0.001 $\pm$ <0.001	0.087 $\pm$ 0.083
	Median Latitude	8.67	1	0.003	0.025 $\pm$ 0.004	0.260 $\pm$ 0.080	
	Region	30.41	3	<0.001	-0.175 $\pm$ 0.027	0.032 $\pm$ 0.038	
	Median Latitude * Historical Distribution Size ($\log_e$)	5.18	1	0.023	-0.001 $\pm$ <0.001	0.581 $\pm$ 0.058	
B	Body Size ($\log_e$)	12.59	1	<0.001	0.529 $\pm$ <0.001	<0.001 $\pm$ <0.001	<0.001 $\pm$ <0.001
	Forest Dependency	50.31	3	<0.001	0.492 $\pm$ <0.001	<0.001 $\pm$ <0.001	
	Residual Generation Time	10.10	1	0.001	0.492 $\pm$ <0.001	0.006 $\pm$ <0.001	
C	Used for Pets	9.15	1	0.002	-1.162 $\pm$ 0.013	<0.001 $\pm$ <0.001	0.269 $\pm$ 0.049
	Used for Food	11.47	1	<0.001	0.353 $\pm$ 0.017	0.029 $\pm$ 0.008	
	Used for Sport	4.55	1	0.033	0.423 $\pm$ 0.019	0.242 $\pm$ 0.021	
D	Residual Human Population Density	3.90	1	0.048	0.196 $\pm$ 0.007	0.006 $\pm$ 0.001	0.324 $\pm$ 0.067
	Residual Urban Population	26.98	1	<0.001	-0.275 $\pm$ 0.013	0.006 $\pm$ 0.002	
	Single Country Endemic	26.44	1	<0.001	0.755 $\pm$ 0.011	<0.001 $\pm$ <0.001	

Table S4. Significant variables in ordinal regression models run separately for each variable group (A-D) predicting the likelihood of a species being more endangered among threatened (VU, EN, CR) parrot species. λ values for the phylogenetic generalized least squares model (PGLS) are given in the table for each model. We give mean $\pm$ standard deviation (SD).

Group	Variable	Deviance	d.f.	$P(\chi^2)$	Estimate $\pm$ SD (PGLS)	P (PGLS) $\pm$ SD	$\lambda \pm$ SD
A	Region	14.42	3	0.002	0.147 $\pm$ <0.001	0.015 $\pm$ <0.001	<0.001 $\pm$ <0.001
	Island Endemic	10.45	1	0.001	-0.250 $\pm$ <0.001	0.210 $\pm$ <0.001	
	Historical Distribution Size (loge)	5.15	1	0.023	-0.016 $\pm$ <0.001	0.621 $\pm$ <0.001	
B	Main Diet	10.04	4	0.040	0.094 $\pm$ <0.001	0.078 $\pm$ <0.001	<0.001 $\pm$ <0.001
C	Used for Pets	4.55	1	0.033	-0.495 $\pm$ <0.001	0.025 $\pm$ <0.001	<0.001 $\pm$ <0.001
D	Per capita GDP	8.79	1	0.003	<0.001 $\pm$ <0.001	0.030 $\pm$ <0.001	<0.001 $\pm$ <0.001
	Single Country Endemic	5.60	1	0.018	0.404 $\pm$ <0.001	0.007 $\pm$ <0.001	

Table S5. (a) Significant variables in the combined linear logistic regression model predicting the likelihood of a parrot species being threatened (VU, EN, CR). λ value for the phylogenetic generalized least squares model (PGLS) was $<0.001 \pm <0.001$. We give mean $\pm$ standard deviation (SD).

Variable	Wald statistic	d.f.	$P(\chi^2)$	Estimate $\pm$ SD (PGLS)	P (PGLS) $\pm$ SD
Historical Distribution Size ($\log_e$)	64.89	1	<0.001	$-0.245 \pm <0.001$	$<0.001 \pm <0.001$
Body Size ($\log_e$)	13.18	1	<0.001	$0.474 \pm <0.001$	$<0.001 \pm <0.001$
Residual Generation Time	16.82	1	<0.001	$0.200 \pm <0.001$	$<0.001 \pm <0.001$
Urban Population	28.71	1	<0.001	$0.014 \pm <0.001$	$<0.001 \pm <0.001$
Forest Dependency	29.47	3	<0.001	$0.332 \pm <0.001$	$<0.001 \pm <0.001$

(b) Significant variables in combined ordinal regression model predicting the likelihood of a species being more endangered among threatened (VU, EN, CR) parrot species. λ value for the final phylogenetic generalized least squares model (PGLS) was $<0.001 \pm <0.001$. We give mean $\pm$ standard deviation (SD).

Variable	Deviance	d.f.	$P(\chi^2)$	Estimate $\pm$ SD (PGLS)	P (PGLS) $\pm$ SD
Per capita GDP	8.47	1	0.004	$0 \pm <0.001$	$0.024 \pm <0.001$
Single Country Endemic	5.60	1	0.018	$0.360 \pm <0.001$	$0.015 \pm <0.001$
Used for Pets	4.75	1	0.029	$-0.526 \pm <0.001$	$0.016 \pm <0.001$

Table S6. (a) Significant threat variables in linear logistic regression model predicting the likelihood of a parrot species being threatened (VU, EN, CR). λ value for the phylogenetic generalized least squares model (PGLS) was 0.009 ± 0.018 .


Variable	Wald statistic	d.f.	$P (\chi^2)$	Estimate $\pm$ SD (PGLS)	P (PGLS) $\pm$ SD
Invasive Alien Species	32.65	1	<0.001	1.734 ± 0.004	$<0.001 \pm <0.001$
Agriculture	28.97	1	<0.001	1.132 ± 0.008	$<0.001 \pm <0.001$
Hunting & Trapping	19.88	1	<0.001	1.383 ± 0.002	$<0.001 \pm <0.001$
Residential & Commercial Development	7.13	1	0.008	0.452 ± 0.009	$<0.001 \pm <0.001$
Energy Production & Mining	4.71	1	0.030	0.397 ± 0.010	0.009 ± 0.002
Agriculture * Hunting & Trapping	7.39	1	0.007	-0.772 ± 0.012	$<0.001 \pm <0.001$
Hunting & Trapping * Invasive Species	6.46	1	0.011	-1.178 ± 0.002	$<0.001 \pm <0.001$

(b) Significant threat variables in ordinal regression model predicting the likelihood of a species being more endangered among threatened (VU, EN, CR) parrot species. λ value for the final phylogenetic generalized least squares model (PGLS) was $<0.001 \pm <0.001$.

Variable	Deviance	d.f.	$P (\chi^2)$	Estimate $\pm$ SD (PGLS)	P (PGLS) $\pm$ SD
Invasive Alien Species	14.41	1	<0.001	$0.598 \pm <0.001$	$<0.001 \pm <0.001$
Agriculture	5.85	1	0.016	$0.332 \pm <0.001$	$0.063 \pm <0.001$

Figure S1. Global density maps of (A) all extant parrot species, and (B) threatened parrots (Vulnerable, Endangered, and Critically Endangered). Color intensities indicate the number of parrot species.

(a)



(b)




Figure S2. Predicted effect of significant variables on the probability of being threatened for parrot species according to: (a) Log_e (Historical Distribution Size, km^2), (b) Log_e (Body Size, cm), (c) Residual Generation Time (years), (d) Forest Dependency, and (e) Percentage of the human population living in urban conditions. Error bars represent standard errors.





Figure S3. Proportion of threatened species in each category (VU, EN, CR) for each significant variable in ordinal regression models: (a) Per capita GDP, (b) Single Country Endemic, (c) whether Used for Pets/Display, and (d) whether threatened by Invasive Alien Species or other threats.



References

BirdLife International (2014) The BirdLife checklist of the birds of the world: Version 7.

Downloaded from

http://www.birdlife.org/datazone/userfiles/file/Species/Taxonomy/BirdLife_Checklist_Version_70.zip

BirdLife International and NatureServe (2014) Bird species distribution maps of the world.

Version 4.0. BirdLife International, Cambridge, UK and NatureServe, Arlington, USA

CIA (2013) Central Intelligence Agency: The World Factbook.

<https://www.cia.gov/library/publications/the-world-factbook/>. Accessed June, 2013

Croxall JP, Butchart SHM, Lascelles B, Stattersfield AJ, Sullivan B, Symes A, Taylor P (2012)

Seabird conservation status, threats and priority actions: a global assessment. Bird Conservation International 22:1-34 doi:10.1017/S0959270912000020

FAOSTAT (2013) Food and Agriculture Organization of the United Nations.

<http://faostat3.fao.org/>. Accessed July, 2013

Hoyo J, Christie DA, Collar NJ, Elliott A, International B, Fishpool LDC, Allen R (2014) HBW

and Birdlife International Illustrated Checklist of the Birds of the World. Lynx Publications, USA

IMF (2013) International Monetary Fund: World Economic Outlook Database.

<http://www.imf.org/>. Accessed June, 2013

IUCN (2014) The IUCN Red List of Threatened Species. Version 2014.2.

<http://www.iucnredlist.org>.

Owens IPF, Bennett PM (2000) Ecological basis of extinction risk in birds: Habitat loss versus human persecution and introduced predators. *Proceedings of the National Academy of Sciences* 97:12144-12148 doi:10.1073/pnas.200223397

Payne RW (2009) *GenStat Wiley Interdisciplinary Reviews: Computational Statistics* 1:255-258 doi:10.1002/wics.32

Salafsky N et al. (2009) Pragmatism and Practice in Classifying Threats: Reply to Balmford et al. *Conservation Biology* 23:488-493 doi:10.1111/j.1523-1739.2009.01197.x

Tittensor DP et al. (2014) A mid-term analysis of progress toward international biodiversity targets. *Science* 346:241-244 doi:10.1126/science.1257484